

the rate of which was lower than reported previously. Overall, our rate of permanent discontinuation of benznidazole was substantially lower than the rates reported previously.^{1,2} We did not measure serum drug concentrations because this would have been costly and impractical in a large randomized trial. We believe that reinfection was unlikely because only 2.6% of the patients lived in an infested house. The mean duration of follow-up in our trial was 5.4 years; however, 75% of the patients were followed for up to 7 years. Although this is a short period as compared with the long natural history of Chagas' disease, it is unknown whether extended follow-up would change our conclusions in the overall population studied.

Cordeiro questions the validity of PCR assay as a marker of therapeutic efficacy or failure. We used it simply to show that benznidazole has an effect on this measure in chronic Chagas' cardiomyopathy, but the effect was indeed more modest than it is in the indeterminate stage of the disease.³ We agree that PCR-based detection of circulating *T. cruzi* is a useful method for denoting treatment failure. However, BENEFIT was a pragmatic trial, and we do not assert that parasite clearance from the blood would not lead to clinical improvement. Our data suggest that the currently used regimen that is recommended by most guidelines^{4,5} may be inadequate. It is possible that in younger patients with no evidence of cardiomyopathy, etiologic treatment to remove or correct the cause of the disease may be more effective, but this determination requires prospective evaluation in larger and longer-term randomized trials.

Urbina et al. raise several issues. PCR conversion rates in the placebo group were higher in our trial than in previously reported studies, but most of the data from those studies were obtained from patients in the indeterminate stage of Chagas' cardiomyopathy and were based on

the inclusion of patients with multiple positive PCR measurements; therefore, those studies cannot be directly compared with ours. With respect to the other issues raised, it is uncertain whether focusing on patients who have a response to treatment as detected by PCR assay would identify a subgroup of the population that would benefit from treatment. Whether newer regimens of trypanocidal drugs, given alone or in combination, or whether repeated dosing or intervening at an earlier stage of the disease would be associated with greater benefit is also uncertain and needs to be explored further. Our study emphasizes the need to test hypotheses, however attractive, in large randomized trials that can provide reliable information on the effects of treatment on clinical outcomes.

Carlos A. Morillo, M.D.

Population Health Research Institute
Hamilton, ON, Canada
morillo@hhsc.ca

Jose A. Marin-Neto, M.D., Ph.D.

University of São Paulo
Ribeirão Preto, Brazil

Alvaro Avezum, M.D., Ph.D.

Instituto Dante Pazzanese de Cardiologia
São Paulo, Brazil

Since publication of their article, the authors report no further potential conflict of interest.

1. Rassi A, Rassi A Jr, Marin-Neto JA. Posaconazole versus benznidazole for chronic Chagas' disease. *N Engl J Med* 2014;371:965.
2. Pérez-Molina JA, Pérez-Ayala A, Moreno S, Fernández-González MC, Zamora J, López-Velez R. Use of benznidazole to treat chronic Chagas' disease: a systematic review with a meta-analysis. *J Antimicrob Chemother* 2009;64:1139-47.
3. Molina I, Gómez i Prat J, Salvador F, et al. Randomized trial of posaconazole and benznidazole for chronic Chagas' disease. *N Engl J Med* 2014;370:1899-908.
4. Andrade JP, Marin Neto JA, Paola AA, et al. I Latin American Guidelines for the diagnosis and treatment of Chagas' heart disease: executive summary. *Arq Bras Cardiol* 2011;96:434-42.
5. Bern C, Montgomery SP, Herwaldt BL, et al. Evaluation and treatment of Chagas disease in the United States: a systematic review. *JAMA* 2007;298:2171-81.

DOI: 10.1056/NEJMc1514453

FTO Obesity Variant and Adipocyte Browning in Humans

TO THE EDITOR: Different polymorphisms in the fat-mass and obesity-associated gene (*FTO*) have been consistently associated with traits related to obesity and were sensibly scrutinized by Claussnitzer et al. (Sept. 3 issue)¹ by means of genetic

and genomic approaches. The authors reported the regulatory importance of the *FTO* locus in early adipocyte differentiation and the causal role of a risk-conferring *FTO* variant (rs1421085) on adiposity, which is believed to act by disrupt-

ing ARID5B-mediated repression of *IRX3* and *IRX5*, thereby influencing mitochondrial respiration and adipocyte browning.

A reduction in adipocyte browning caused by a developmental swing between energy-dissipating beige (brite) fat cells and energy-accumulating white adipocytes reduces thermogenesis and subsequently increases lipid deposition, which in turn supports investigations that have reported uncoupling functions in association with *FTO* and *UCP1*.² However, the involvement of other recognized *FTO* polymorphisms that affect fuel utilization through appetite or the control of macronutrient intake³ or through the sensing of amino acids and nutrients,⁴ and the potential epigenetic implications of *FTO* as a DNA–RNA demethylase-modifying enzyme or an epitranscriptomic marker,⁵ should not be disregarded. Furthermore, causal or consequential reanalyses of the interactive or overlapping roles of *FTO* in energy metabolism need additional research.

Fermin I. Milagro, Ph.D.

Maria J. Moreno-Aliaga, Ph.D.

J. Alfredo Martinez, M.D., Ph.D.

University of Navarra
Pamplona, Spain
jalfmtz@unav.es

No potential conflict of interest relevant to this letter was reported.

1. Claussnitzer M, Dankel SN, Kim KH, et al. *FTO* obesity variant circuitry and adipocyte browning in humans. *N Engl J Med* 2015;373:895-907.
2. Tews D, Fischer-Posovszky P, Fromme T, et al. *FTO* deficiency induces *UCP-1* expression and mitochondrial uncoupling in adipocytes. *Endocrinology* 2013;154:3141-51.
3. Livingstone KM, Celis-Morales C, Lara J, et al. Associations between *FTO* genotype and total energy and macronutrient intake in adults: a systematic review and meta-analysis. *Obes Rev* 2015;16:666-78.
4. Gulati P, Yeo GS. The biology of *FTO*: from nucleic acid demethylase to amino acid sensor. *Diabetologia* 2013;56:2113-21.
5. Zhao X, Yang Y, Sun BF, Zhao YL, Yang YG. *FTO* and obesity: mechanisms of association. *Curr Diab Rep* 2014;14:486.

DOI: 10.1056/NEJMc1513316

TO THE EDITOR: Claussnitzer et al. examined the mechanisms through which *FTO* variants are associated with fatness and conclude that these variants reduce thermogenesis in human adipocytes. However, the authors did not determine whether carriers of alleles associated with an increased risk of obesity have reduced energy expenditure. Speakman¹ reviewed 14 published studies and concluded that there is no evidence of such an effect. In contrast, several of the stud-

ies reviewed by Speakman reported that carriers of the *FTO* risk variants had alterations in appetite, which is consistent with a growing body of evidence indicating that the expression of genes implicated in human fatness is markedly enriched in the nervous system.² Tschöp et al.³ have recommended methods for the comparison of energy expenditures in mice with different body compositions, but these methods were not adopted by Claussnitzer et al., which reduces confidence in their results. Cellular and murine models can be powerful tools when used to dissect the mechanisms underlying the association of genetic variants with human phenotypes. However, these models are not a substitute for the detailed phenotyping of the species in which the association was discovered.

Stephen O'Rahilly, M.D.

Anthony P. Coll, M.B., B.S., Ph.D.

Giles S.H. Yeo, Ph.D.

University of Cambridge
Cambridge, United Kingdom
so104@medschl.cam.ac.uk

No potential conflict of interest relevant to this letter was reported.

1. Speakman JR. The 'fat mass and obesity related' (*FTO*) gene: mechanisms of impact on obesity and energy balance. *Curr Obesity Reports* 2015;4:73-91.
2. Locke AE, Kahali B, Berndt SI, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015; 518:197-206.
3. Tschöp MH, Speakman JR, Arch JR, et al. A guide to analysis of mouse energy metabolism. *Nat Methods* 2012;9:57-63.

DOI: 10.1056/NEJMc1513316

TO THE EDITOR: Claussnitzer et al. unravel the switchlike behavior of the single *FTO* nucleotide variant rs1421085 in white adipocyte browning and thermogenesis through an evolutionarily conserved motif for the ARID5B repressor that regulates the downstream expression of *IRX3* and *IRX5*. Yet the idea that mitochondrial thermogenesis is regulated through *FTO* is not new. The authors did not cite a seminal discovery that links *FTO* deficiency to an induction of beige adipocytes by means of *UCP1* expression and mitochondrial uncoupling, as reported by Tews et al. in 2013.¹ Moreover, they did not consider an epitranscriptomic level of regulation of target-gene expression critical to adipocyte browning, despite the fact that *FTO* was recently described as encoding an RNA demethylase in humans.² Analogous to DNA methylation and histone modification in epigenomics, *FTO* influences differential

adenosine methylation in messenger RNA on a transcriptome-wide scale³ and alters RNA splicing, stability, and protein binding.⁴ *FTO* deficiency has led to an increase in the expression of *UCP1* by a factor of 4 and of *PGC-1 α* by a factor of 2, with correspondingly higher uncoupled mitochondrial respiration rates. RNA sequencing is warranted to elucidate the role of N⁶-methyladenosine epitranscriptomics in the molecular and biologic phenomena related to *FTO*-mediated browning and mitochondrial thermogenesis, as observed above.

Melvin K. Leow, M.D., Ph.D.

Brenner Center for Molecular Medicine
Singapore, Singapore
mleowsj@massmed.org

No potential conflict of interest relevant to this letter was reported.


1. Tews D, Fischer-Posovszky P, Fromme T, et al. *FTO* deficiency induces UCP-1 expression and mitochondrial uncoupling in adipocytes. *Endocrinology* 2013;154:3141-51.
2. Jia G, Fu Y, Zhao X, et al. N⁶-methyladenosine in nuclear RNA is a major substrate of the obesity-associated *FTO*. *Nat Chem Biol* 2011;7:885-7.
3. Berulava T, Ziehe M, Klein-Hitpass L, et al. *FTO* levels affect RNA modification and the transcriptome. *Eur J Hum Genet* 2013;21:317-23.
4. Liu N, Dai Q, Zheng G, He C, Parisien M, Pan TN. N(6)-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions. *Nature* 2015;518:560-4.

DOI: 10.1056/NEJMc1513316

THE AUTHORS REPLY: We agree that additional mechanisms in the *FTO* locus may contribute to obesity, including some that act through appetite. However, given the extensive feedback involved in the regulation of energy balance, differences in adult nutrition may result from the differences in adipocyte browning elucidated in our study. For example, a longitudinal study¹ involving 438 children concluded that *FTO* risk variants were significantly associated with body-mass index (BMI) but not with energy intake or physical activity at 15 years of age, indicating that nutritional changes in adults may occur downstream of changes in BMI during adolescence.

We agree with O’Rahilly et al. that detailed human phenotyping is important. However, the studies reviewed by Speakman² are associational, not mechanistic, and thus cannot determine whether differences in energy intake are a cause or a consequence of differences in body fat. The cited studies are also underpowered to attribute different subsets of phenotypic associations

to different subsets of variants, given the high linkage disequilibrium in the locus ($D' > 0.99$, $r^2 > 0.93$). Moreover, the studies cited by Speakman reported only resting energy expenditure — not stimulus-induced energy expenditure (e.g.,



by cold or exercise) — for which we found the strongest genetic effect in our study.

As noted by Leow, several authors, including Tews et al.,³ argue that the *FTO* locus acts on obesity through the m6A epitranscriptomic function of the *FTO* protein. However, only *IRX3* and *IRX5* showed a significant difference in their levels of adipocyte expression between risk-allele carriers and nonrisk allele carriers. Moreover, several manipulations described by Tews et al. may have altered *IRX3* and *IRX5* expression levels (which were not reported), possibly explaining the resulting phenotypes. Last, patients who have homozygous loss-of-function mutations affecting the *FTO* protein have severe growth retardation and multiple malformations but no effect of obesity,⁴ indicating the *FTO* protein is unlikely to be a primary phenotypic mediator, irrespective of the tissue of action.

As suggested by O’Rahilly et al., we also use analysis of covariance (ANCOVA) to compare the energy expenditure of *Irx3* dominant-negative (aP2-*Irx3*DN) and control mice, with adjustment for lean body mass with the use of ANCOVA. We continue to find significant differences in energy expenditure (Fig. 1), which support our initial conclusions. Last, we agree that additional studies are needed to continue the process of eluci-

dating the ways in which genetic variants in the *FTO* locus affect obesity beyond the ARID5B–rs1421085–*IRX3/IRX5*–browning axis elucidated in our study.

Melina Claussnitzer, Ph.D.

Harvard Medical School
Boston, MA
melina@broadinstitute.org

Chi-Chung Hui, Ph.D.

Hospital for Sick Children
Toronto, ON, Canada

Manolis Kellis, Ph.D.

Massachusetts Institute of Technology
Cambridge, MA

Since publication of their article, the authors report no further potential conflict of interest.

1. Hakanen M, Raitakari OT, Lehtimäki T, et al. *FTO* genotype is associated with body mass index after the age of seven years but not with energy intake or leisure-time physical activity. *J Clin Endocrinol Metab* 2009;94:1281-7.
2. Speakman JR. The ‘fat mass and obesity related’ (*FTO*) gene: mechanisms of impact on obesity and energy balance. *Curr Obesity Reports* 2015;4:73-91.
3. Tews D, Fischer-Posovszky P, Fromme T, et al. *FTO* deficiency induces UCP-1 expression and mitochondrial uncoupling in adipocytes. *Endocrinology* 2013;154:3141-51.
4. Boissel S, Reish O, Proulx K, et al. Loss-of-function mutation in the dioxygenase-encoding *FTO* gene causes severe growth retardation and multiple malformations. *Am J Hum Genet* 2009;85:106-11.

DOI: 10.1056/NEJMc1513316

Chimeric Antigen Receptor T Cells in Myeloma

TO THE EDITOR: Garfall et al. (Sept. 10 issue)¹ describe an interesting case of a patient with multiple myeloma who was successfully treated with autologous stem-cell transplantation followed by infusion of anti-CD19 chimeric antigen receptor–transduced CTL019 cells. The vast majority (99.95%) of the patient’s myeloma cells did not express CD19; this indicates that the effects of CTL019 could not be attributed to direct cytotoxicity against the predominant malignant clone itself.

The authors hypothesize that the therapeutic activity of CTL019 is due to targeting of rare CD19-expressing myeloma precursors. However, an alternative explanation for the antimyeloma effect of CTL019 therapy could be that CTL019 results in depletion of nonmalignant CD19-positive B cells. Studies have shown that B cells can promote cancer by providing growth and survival

factors for the tumor and by suppressing antitumor immunity.²⁻⁴ Of note, patients with multiple myeloma frequently have altered B-cell homeostasis with a characteristic expansion of a distinct population of memory B cells.⁵ Thus, CTL019-mediated depletion of a pro-myeloma B-cell population could contribute to the clinical benefit of CTL019.

Alexander Shimabukuro-Vornhagen, M.D.

Hans A. Schloesser, M.D.

Michael S. von Bergwelt-Baildon, M.D., Ph.D.

University Hospital of Cologne
Cologne, Germany
shima@uk-koeln.de

No potential conflict of interest relevant to this letter was reported.

1. Garfall AL, Maus MV, Hwang W-T, et al. Chimeric antigen receptor T cells against CD19 for multiple myeloma. *N Engl J Med* 2015;373:1040-7.
2. de Visser KE, Korets LV, Coussens LM. De novo carcinogen-