



Genetic drivers of m⁶A methylation in human brain, lung, heart and muscle

Xushen Xiong^{1,2,3,4}, Lei Hou^{1,2,4}, Yongjin P. Park^{1,2}, Benoit Molinie², GTEx Consortium*, Richard I. Gregory³ and Manolis Kellis^{1,2} ✉

The most prevalent post-transcriptional mRNA modification, N⁶-methyladenosine (m⁶A), plays diverse RNA-regulatory roles, but its genetic control in human tissues remains uncharted. Here we report 129 transcriptome-wide m⁶A profiles, covering 91 individuals and 4 tissues (brain, lung, muscle and heart) from GTEx/eGTEx. We integrate these with interindividual genetic and expression variation, revealing 8,843 tissue-specific and 469 tissue-shared m⁶A quantitative trait loci (QTLs), which are modestly enriched in, but mostly orthogonal to, expression QTLs. We integrate m⁶A QTLs with disease genetics, identifying 184 GWAS-colocalized m⁶A QTL, including brain m⁶A QTLs underlying neuroticism, depression, schizophrenia and anxiety; lung m⁶A QTLs underlying expiratory flow and asthma; and muscle/heart m⁶A QTLs underlying coronary artery disease. Last, we predict novel m⁶A regulators that show preferential binding in m⁶A QTLs, protein interactions with known m⁶A regulators and expression correlation with the m⁶A levels of their targets. Our results provide important insights and resources for understanding both *cis* and *trans* regulation of epitranscriptomic modifications, their interindividual variation and their roles in human disease.

Genome-wide association studies (GWAS) have identified >100,000 genetic loci associated with complex traits and diseases, but 93% do not affect protein-coding regions directly and remain largely uninterpreted, hindering efforts to decipher the molecular basis of human disease^{1,2}. To bridge this gap between genetic variation and disease phenotypes, many studies have profiled diverse molecular processes affected by non-coding variants, revealing thousands of genetic variants impacting gene expression (expression QTLs (eQTLs)), mRNA splicing, DNA methylation, histone acetylation, RNA editing and protein levels^{3–11}, and providing insights into increasing numbers of GWAS loci. However, the role of genetic variants affecting post-transcriptional mRNA modifications in human tissues and complex diseases remains uncharacterized.

More than 150 types of post-transcriptional RNA modifications regulate non-coding RNAs and protein-coding mRNA transcripts, similarly to epigenomic modifications of DNA and histone modifications^{12,13}. Among the many such mRNA ‘epitranscriptomic’ marks, m⁶A is the most prevalent, and most frequently associated with development and disease^{12,14–18}. m⁶A modulates mRNA maturation, degradation, export and translation efficiency, and is involved in spermatogenesis, stem cell differentiation, immune response, neurodevelopment, tumorigenesis and other processes^{14,16,17,19–21}. It is deposited by m⁶A ‘writer’ methyltransferases (METTL3 and METTL14) and adapters (WTAP, VIRMA, ZC3H13, CBL1, RBM15 and RBM15B)^{12,22} at preferential sequence motifs¹², removed by m⁶A ‘eraser’ demethylases (FTO, also known as ALKBH9, and ALKBH5) and recognized by m⁶A ‘readers’ (YT521-B homology (YTH) family) and other direct or indirect binders (IGF2BP1–3, FMR1, HNRNPC/G and HNRNPA2B1), with different ‘readers’ leading to decay (for example, YTHDF2) or stabilization (IGF2BP proteins). However, many m⁶A regulators remain uncharacterized^{23–25}.

Diverse diseases are linked to m⁶A dysregulation, including glioblastoma, lung cancer and acute myeloid leukemia^{16,20}, and m⁶A

loci are enriched in disease-associated variants, possibly indicating causal roles^{26,27}. Genetic loci controlling the m⁶A level as a quantitative trait (m⁶A QTLs) were identified in lymphoblastoid cell lines (LCLs) derived from 60 individuals of YRI ancestry (Yoruba in Ibadan, Nigeria)²⁸, but m⁶A QTLs remain uncharacterized in primary human tissues or in other genetic backgrounds.

Here we report a collection of transcriptome-wide post-transcriptional modifications in human primary tissues, spanning 176 samples across 107 individuals in brain, lung, heart and muscle. We discover 9,312 m⁶A QTLs in brain, lung, heart and muscle targeting 1,270 m⁶A sites (genetically driven m⁶A (g-m⁶A) sites). Most m⁶A QTLs are distinct from eQTLs, but the small subset of eQTL-colocalized m⁶A QTLs are preferentially degradation-associated. m⁶A QTLs enrich in disease-associated loci. Several m⁶A QTLs help elucidate GWAS loci, including brain m⁶A QTLs for psychiatric disorders, lung m⁶A QTLs for expiratory flow and asthma, and muscle/heart m⁶A QTLs for heart disease and high blood pressure. We predict new m⁶A-regulating RNA-binding proteins (RBPs) with preferential binding in m⁶A-QTL single-nucleotide polymorphisms (SNPs), experimentally confirmed interactions with known m⁶A regulators and significant expression correlation with m⁶A levels of their targets.

Results

m⁶A variation across tissues and individuals. We generated 176 transcriptome-wide m⁶A profiles across 107 individuals in human brain, lung, heart and muscle (Supplementary Table 1), as part of the enhancing GTEx (eGTEx) consortium²⁹. We selected individuals from the Genotype-Tissue Expression (GTEx) project^{4,8,30}, enabling us to directly compare our discovered m⁶A QTLs with mRNA expression eQTLs and other pre- and post-transcriptional processes.

We extended methylated-RNA immunoprecipitation with sequencing (meRIP-seq)^{18,31,32} protocols for m⁶A identification and

¹Computer Science and Artificial Intelligence Lab, Massachusetts Institute of Technology, Cambridge, MA, USA. ²The Broad Institute of Harvard and MIT, Cambridge, MA, USA. ³Stem Cell Program, Division of Hematology/Oncology, Boston Children’s Hospital, Boston, MA, USA. ⁴These authors contributed equally: Xushen Xiong, Lei Hou. *A list of authors appears at the end of the paper. ✉e-mail: manoli@mit.edu

quantification, which we optimized for 400-fold-reduced starting material from GTEx compared to previous studies³¹, and used matched RNA sequencing (RNA-seq) data as the background (Fig. 1a). Our aggregated meRIP-seq signal showed strong enrichment surrounding the known GGACH m⁶A deposition sequence motif (centered on the modified A base; Extended Data Fig. 1a). After stringent quality controls using peak calling, positional enrichment and consensus motif enrichment, we focused on a subset of 129 high-quality m⁶A maps across 91 individuals that also have genetic information, including 53 brain, 12 heart, 32 muscle and 32 lung samples (Fig. 1a and Extended Data Fig. 1a–c).

Across all samples, we found >278,000 multiply detected m⁶A sites (detected in ≥2 individuals), with an average of ~20,000 sites per sample (Extended Data Fig. 1d). Our profiling greatly expanded the previously known set of m⁶A methylated sites, with 77% of our multiply detected sites in brain, 60% in heart, 58% in muscle and 72% in lung not previously reported³³ (Extended Data Fig. 1e). Our m⁶A sites were enriched for the known m⁶A consensus motif, and showed enrichment near stop codons, consistent with previous studies^{31,32,34} (Extended Data Fig. 1b,c). Previously undetected m⁶A sites showed positional (Extended Data Fig. 1f) and motif enrichment (adjusted $P = 5.7 \times 10^{-26}$, Fisher test) equally as strong as that for previously detected sites, confirming their high quality.

Clustering all m⁶A profiles by their global similarity, we found that the tissue profiled was the primary driver of variation (Fig. 1b), with brain samples most distinct, and heart and muscle clustering first together and then with lung, consistent with tissue-specific biology^{26,27,35} and with RNA-based tissue similarity (Extended Data Fig. 1g). Genes harboring tissue-specific m⁶A sites showed tissue-relevant functional enrichments (Extended Data Fig. 1h), including synaptic and neuronal pathways in brain, and cardiomyopathy and muscle-related functions in muscle and heart. Noteworthy examples of tissue-specific methylation include brain-specific m⁶A in *POU3F2* (Fig. 1c), where m⁶A regulates glioblastoma³⁶, and lung-specific m⁶A for *EGFR* (Fig. 1d), where m⁶A promotes lung cancer³⁵. While genes with tissue-shared m⁶A sites show little expression difference across tissues (Fig. 1e), tissue-specific m⁶A sites do not primary stem from mRNA tissue-specific expression: for 72.8% of tissue-specific m⁶A cases, the corresponding transcript was broadly expressed in the other tissues but m⁶A modifications were not detected in those tissues (Fig. 1f), indicating that m⁶A can lead to tissue-specific functions for otherwise broadly expressed transcripts.

m⁶A genetic driver discovery and validation. We next sought to recognize m⁶A QTLs and their target m⁶A (g-m⁶A) sites. Given the smaller sample size in heart, we called m⁶A QTLs in heart and muscle jointly to gain power, as heart and muscle were previously shown to share many eQTLs⁸, and co-cluster for both GTEx mRNA data (Extended Data Fig. 1g) and our m⁶A data (Fig. 1b).

To recognize and remove ‘unwanted’ global variation that confounds and hides subtle effects of *cis*-acting genetic variants^{8,37}, we used a tool based on a Bayesian factor (PEER³⁷). We removed the top seven factors in brain, the top five in muscle/heart and the top two in lung, as inclusion of additional factors led to a rapid decrease in the number of g-m⁶A sites (Fig. 2a), and additional factors were highly correlated with each other, indicating signal saturation (Fig. 2b and Extended Data Fig. 2a). The removed factors correlated with batch, sex, age and ancestry, as expected, and also immune disease, psychiatric traits and medication history (Fig. 2c and Extended Data Fig. 2b), which are expected confounders acting in non-local ways whose removal decreases noise during QTL analyses.

We used FastQTL to perform a permutation-based m⁶A-QTL search for each m⁶A site, calculating the empirical *P* value for the SNP with the strongest genetic effect for each m⁶A site. As m⁶A modifications are deposited both co-transcriptionally and

post-transcriptionally, we searched for m⁶A-SNPs in promoter regions (where they can act at the DNA level), in introns (where they can act both at the DNA level and at the RNA level before splicing) and in exons (including UTRs and coding regions, where they act at the DNA or RNA level, either pre- or post-splicing).

This resulted in 9,312 m⁶A QTLs targeting 1,058 genes across the 4 tissues, of which ~5,200 m⁶A QTLs act in brain (472 genes), ~3,100 in lung (401 genes) and ~2,100 in muscle/heart (279 genes), after correcting for multiple-SNP testing in each locus (empirical $P < 0.005$; (Fig. 2d, Extended Data Fig. 2c,d and Supplementary Table 2). The higher sample size in brain allowed us to also report ~1,300 higher-confidence brain m⁶A QTLs (targeting 94 genes; Supplementary Table 3), using 2 rounds of multiple-testing correction for both multiple SNPs and multiple m⁶A targets (see Methods). These m⁶A QTLs showed substantial genotype-driven differences in m⁶A levels between individuals, and often affected m⁶A levels of biologically important genes in relevant tissues, including the neuronal function genes *PADI2*, *MOBP* and *DLG1* in brain, the lung function and respiratory genes *CFLAR*, *SPTBN1* and *SLIT2* in lung, and the heart/skeletal muscle function genes *PDIA6*, *SSRI* and *HSPA8* in muscle/heart (Fig. 2e and Extended Data Fig. 2c,d).

To validate our m⁶A-QTL results (in our discovery study), we used the m⁶A levels measured in muscle, heart and lung from two individuals²⁶ (validation study). As two individuals are insufficient for m⁶A-QTL discovery, we tested whether the effect size direction matched between the two studies, focusing on $n = 281$ m⁶A-QTL SNPs (5.6%) for which the two individuals from the validation study have different genotypes, and separating our m⁶A QTLs into positive effect (minor allele shows a significantly higher m⁶A level), negative effect (minor allele significantly lower) and no effect (no significant m⁶A difference). Despite using only two individuals, and despite potential population-specific effects (between our European-ancestry discovery study and the Asian-ancestry validation) that may reduce agreement, we found a consistent and significant agreement between our discovery study and the validation study (Fig. 2f): our positive-effect m⁶A QTLs showed a 28% higher median m⁶A level in the validation study ($P = 0.04$, Wilcoxon test), our negative-effect m⁶A QTLs showed a 36% lower median m⁶A level in validation ($P = 5.5 \times 10^{-5}$, Wilcoxon test), and our no-effect SNPs showed only a 1.007-fold difference between alleles (used as the baseline to compute the aforementioned *P* values).

Tissue specificity of m⁶A QTLs. Only 5.0% of m⁶A QTLs (6.8% of g-m⁶A sites) were shared between any pair of tissues (Fig. 3a,b and Extended Data Fig. 3a), even though 46% of m⁶A peaks were shared (Fig. 1e), with tissue-specific m⁶A QTLs enriched in distinct functional pathways (Extended Data Fig. 3b), including synaptic function and signaling in brain.

To evaluate whether m⁶A-QTL specificity stemmed from biological factors or lack of discovery power, we tested whether the 8,800 tissue-specific m⁶A QTLs (~95% of all m⁶A QTLs) showed consistent directionality in the tissues where they were not discovered as m⁶A QTLs. We reasoned that if the m⁶A-QTL effects were shared in those tissues, we would see common directionality, despite having limited power to detect them, with complete sharing resulting in 100% effect directionality agreement, and complete tissue specificity resulting in 50% effect directionality agreement expected by chance. Indeed, we found a gradual transition from 100% consistency to near-50% consistency (Fig. 3c,d), at decreasing *P*-value thresholds for the second tissue: the 469 shared m⁶A QTLs (5%) that were significant in both tissues showed 100% directionality consistency, providing strong confirmation of our results based on independent discovery in multiple tissues (red in Fig. 3d); an additional 592 m⁶A QTLs (6.4%) that were significant in one tissue ($P < 0.005$) and near-threshold *P* values in another tissue ($0.005 < P < 0.05$) showed 99% consistency, indicating that our current threshold




Fig. 1 | Study design and m⁶A landscape across tissues. **a**, An overview of the m⁶A profiling method and samples. Created with BioRender.com. **b**, Pearson correlation (heatmap) and hierarchical clustering (tree) of m⁶A profiles across 129 samples (rows/columns) shows tissue-specific clustering (colors). **c**, An example of brain-specific m⁶A (top) in the *POU3F2* 3'-UTR with brain-specific RNA-seq expression (bottom). **d**, An example of lung-specific m⁶A (top) in the *EGFR* 3'-UTR despite tissue-shared expression (bottom). **e**, Tissue-shared m⁶A (top) and corresponding expression (bottom) across peaks (columns) and tissues (rows). **f**, Tissue-specific m⁶A (top) and corresponding expression (bottom) across peaks (columns) and tissues (rows), showing that 72.8% of tissue-specific m⁶A sites lack tissue-specific expression (for example, **d**) and 27.2% also show tissue-specific expression (for example, **c**).

for m⁶A-QTL discovery is quite stringent, and that even looser thresholds for m⁶A-QTL discovery would still result in additional meaningful m⁶A QTLs; another ~1,680 m⁶A QTLs (~18%) that were significant in one tissue and subthreshold in the second tissue ($0.05 < P < 0.5$) showed 92% consistency, indicating increasing numbers of false positives at those thresholds; the remaining ~6,600 m⁶A QTLs (~71%) that were significant in one tissue and showed

$P > 0.5$ in the second tissue showed only 60% consistency in m⁶A genetic effect directionality (near the 50% expected by chance), indicating that the vast majority of calls would be false positives at such thresholds.

Leveraging the fact that our m⁶A profiling was performed in individuals from the GTEx cohort, we also directly compared the tissue sharing of eQTLs and the tissue sharing of m⁶A QTLs in the exact




Fig. 2 | Genetically driven m⁶A sites across tissues. **a**, The number of g-m⁶A sites (nominal $P < 10^{-4}$; the P value was calculated by FastQTL linear regression) identified (y axis) versus the number of PEER factors used (x axis) shows the inflection point for each tissue (colors). **b**, Pearson correlation (color) for PEER factors in brain shows saturation (correlated factors) after the first seven factors (red box). **c**, Pearson correlation between PEER factors (columns) and donor phenotypic measurements (rows) helps interpret factors. **d, e**, Manhattan plot showing the genomic position (x axis) and the nominal P value for association (y axis) for all m⁶A QTLs in brain (**d**), highlighting three examples (**e**). The box shows the 25th–75th percentile (that is, the interquartile range; IQR); the line shows the median; the whiskers show $1.5 \times$ IQR. The P values were calculated by FastQTL linear regression. **f**, The minor allele effect size in the validation cohort (y axis) for increased-effect m⁶A QTLs (+) and decreased-effect m⁶A QTLs (–) relative to non-m⁶A-QTLs (SNPs tested but not significant during m⁶A-QTL calling, ctrl). The data are from Liu et al.²⁶, and merged m⁶A profiles from two samples of muscle, lung and heart, each. The P values were calculated using a two-sided Wilcoxon test. The box shows the 25th–75th percentile; the line shows the median; the whiskers show $1.5 \times$ IQR.

same cohort of individuals, by subsampling the transcriptional data from GTEx to the specific individuals studied here, thus matching discovery power, allele frequencies and potential *trans*-acting or environmental effects. In the most stringent eQTL threshold (two-step correction, false discovery rate (FDR)=0.1), we found that 14% of eQTLs were independently discovered in multiple tissues (Extended Data Fig. 3c), substantially more than the 5% sharing found for m⁶A QTLs, suggesting that m⁶A QTLs are approximately threefold more tissue specific than eQTLs for matched cohorts. Repeating our directionality consistency analysis for eQTLs across both the full GTEx cohort and a sample-size-matched subsampled GTEx dataset, we found that directionality consistency is robust to sample size (90% for full dataset versus 89% for subsampled eQTLs; Extended Data Fig. 3d,e), and is substantially higher for eQTLs than for m⁶A QTLs (89% for eQTLs versus 71% for m⁶A QTLs, compared to 50% expected by chance; Extended Data Fig. 3e and Fig. 3d), indicative of approximately twice as much eQTL tissue sharing than m⁶A-QTL tissue sharing. We also used simulations with matching effect size, minor allele frequency (MAF) and sample size under the assumption of 100% tissue sharing, and found much higher overlap and directionality consistency between tissues than observed in our measurements (38–90% versus ~5%; Extended Data Fig. 3f,i), indicating that the observed tissue specificity does not simply stem from the discovery power.

We next asked whether the tissue specificity of m⁶A QTLs stems from strong differences in gene expression levels. We first evaluated whether genes with tissue-specific m⁶A QTLs in a given tissue show higher relative expression levels than genes with tissue-shared m⁶A QTLs in the same tissue (Extended Data Fig. 3j), and found that in all cases there was no substantial difference in the expression level between tissue-specific and tissue-shared m⁶A QTLs. We next

evaluated whether genes with tissue-specific m⁶A QTLs in a given tissue show higher expression levels in that tissue than in tissues where they lack m⁶A QTLs (Extended Data Fig. 3k), and found that in four out of six pairwise comparisons, the differences in expression are not significant, and in the last two comparisons, gene expression levels are within 50% of each other. These results indicate that the observed m⁶A-QTL tissue specificity does not simply stem from differences in expression levels.

We next asked whether the primary tissue m⁶A QTLs discovered here were already captured in the m⁶A QTLs previously reported²⁸ in immortalized LCLs, derived from peripheral B lymphocytes by Epstein–Barr virus transformation in 60 Geuvadis YRI samples^{30,38}. As a benchmark, we first evaluated the sharing of eQTLs from the 4 primary tissues profiled here in the full GTEx cohort (205 Brain, 515 Lung, 386 Heart and 706 Muscle samples) and eQTLs from YRI LCLs in the full Geuvadis cohort (89 LCL samples)^{30,38}. This comparison showed strong eQTL sharing between primary tissues of the GTEx samples ($R^2 = 0.66–0.80$) but very low sharing with YRI LCLs ($R^2 = 0.06–0.12$; Extended Data Fig. 3l), as expected given the substantially different biology of immortalized cell lines and primary tissues, compounded with potential differences between EUR and YRI ancestry groups, and indicating that the two capture different parts of biological diversity. Consistent with these eQTL results, the findings for our m⁶A QTLs in primarily EUR-ancestry tissues showed them to be largely distinct from m⁶A QTLs of YRI-ancestry immortalized cell lines, with only 0.8% shared m⁶A QTLs (Extended Data Fig. 3m), very low correlation ($R^2 = 0.0007–0.021$; Extended Data Fig. 3n) and only 51.25% directionality agreement (Extended Data Fig. 3o), which is close to the 50% expected by chance ($P = 0.31$) and substantially lower than the 71% consistency found between tissues ($P < 2.2 \times 10^{-16}$). However, the 0.8% of m⁶A




Fig. 3 | m⁶A-QTL tissue specificity. **a**, m⁶A-QTLs are highly tissue specific. **b**, Tissue specificity of m⁶A-QTL target peaks. Among m⁶A peaks that are m⁶A-QTL targets (that is, g-m⁶A genes) 12–32% are tissue-specific peaks (green), but even among tissue-shared peaks, only 5.0–6.3% have shared m⁶A-QTLs (red). **c**, m⁶A-QTL effect size consistency. Tissue-shared m⁶A-QTLs (red triangles) show 100% consistent effect sizes between tissues (top-right and bottom-left quadrants), and even tissue-specific m⁶A-QTLs (circles) that are near-threshold in a second tissue (red shading) show highly consistent effects (99% agreement for $P < 10^{-2}$) for all pairs of tissues (9 panels). The P value was calculated by FastQTL linear regression. **d**, Between-tissue consistency (right y axis) of the m⁶A-QTL effect size (left y axis) increases with the significance of the tissue-2 m⁶A-QTL P value (x axis) for both positive-effect tissue-1 m⁶A-QTLs (right half-plane) and negative-effect tissue-1 m⁶A-QTLs (left half-plane). The top histogram shows the distribution of cases in each P value; most m⁶A-QTLs in tissue 1 show no effect in the second tissue (the gray bar in the middle). The percentage of loci in each group and the corresponding directionality consistency are shown in parentheses. The P value was calculated by FastQTL linear regression.

QTLs that were shared ($n = 26$) showed 100% consistent directionality (green dots in Extended Data Fig. 3n), providing a form of additional validation for the validity of both studies. These results highlight the importance of also profiling m⁶A-QTLs in primary human tissues of multiple ancestry backgrounds, as they capture additional biological information not captured in immortalized cell lines from a single ancestry background.

m⁶A-QTLs and eQTLs sometimes overlap but are mostly independent. As m⁶A can increase mRNA stability (for example, via IGF2BP)³⁹ or degradation (for example, via YTHDF2)⁴⁰, m⁶A-QTL effects may be positively or negatively correlated with m⁶A target gene expression. Across all SNPs, m⁶A-QTLs showed a mild but significant enrichment in GTEx matched-tissue eQTLs, across brain (1.5-fold, $P = 8.3 \times 10^{-7}$, prefrontal cortex BA9 eQTLs), muscle/heart (2-fold, $P = 0.0026$, muscle eQTLs; 1.6-fold, $P = 0.0050$, left ventricle eQTLs) and lung (1.6-fold, $P = 0.0030$, lung eQTLs; Fig. 4a). Fold-enrichments were similar when exonic and intronic QTLs were considered separately, although significance levels were lower due to the lower number of data points (Extended Data Fig. 4a,b).

Among m⁶A-QTLs colocalized with eQTLs in matched individuals (after PEER factor removal; Extended Data Fig. 4c), 23% were positively correlated with gene expression, reflecting stability-increasing effects (Fig. 4b,c), and 77% were negatively correlated, reflecting degradation effects (Fig. 4b,c). Repressive-effect m⁶A-QTLs were significantly higher (odds ratio = 2.5, $P = 2.3 \times 10^{-14}$, Fisher test) indicating more widespread degradation effects in the tissues studied here. Degradation-mediating g-m⁶A sites preferentially localized in CDS and 3'-UTRs, and stabilization-mediating g-m⁶A sites preferentially localized in long non-coding RNAs and 5'-UTRs, although these differences were not significant due to small counts (Extended Data Fig. 4d).

Despite these enrichments, 88% of m⁶A-QTLs showed no significant eQTL effect and 94% of eQTLs showed no significant

m⁶A-QTL effect in matched samples (no P value < 0.005 , no effect size > 0.5 ; Fig. 4c,d), and m⁶A-QTL target genes (gmGenes, for genetically controlled m⁶A-level genes) and eQTL target genes (eGenes) showed very little overlap (Extended Data Fig. 4e; $P = 0.21$), with 93% of gmGenes lacking eQTLs and 96% of eGenes lacking m⁶A-QTLs, indicating that m⁶A-QTL effects are not simply a consequence of changes in expression. Even across the full GTEx cohort for matching tissues, 79% of m⁶A-QTLs do not show any eQTL effect (Extended Data Fig. 4f), and conversely 99% of eQTLs do not show any m⁶A-QTL effect (Extended Data Fig. 4g). Even when a gmGene was also an eGene, their lead QTL SNPs differed greatly (~10-kilobase (kb) average distance; Fig. 4e), indicating that m⁶A-QTLs and eQTLs constitute largely independent gene-regulatory paths.

m⁶A-QTLs help interpret GWAS loci. Given the known roles of m⁶A in multiple human diseases^{16,17,20,21}, including cancer, immune disease and brain disorders, we next asked whether our discovered m⁶A-QTLs can help shed light on potential mechanisms for currently unexplained genome-wide significant and subthreshold GWAS loci, and possibly implicate m⁶A function in additional disorders.

We found 179 m⁶A-QTLs colocalized with 80 GWAS traits (Supplementary Table 4), with 88% ($n = 157$) of lead SNPs outside protein-coding exons, and 82% ($n = 147$) distinct from eQTLs in the full GTEx cohort for our tissues, thus providing new tissue-specific mechanistic hypotheses for these loci, including candidate causal variants, target genes and tissues of action.

For brain, 71 m⁶A-QTLs colocalize with GWAS variants. Neuroticism-associated rs12471193 is a brain m⁶A-QTL for the potassium channel gene *KCNJ3*, a neuronal excitability regulator implicated in multiple brain disorders⁴¹ (Fig. 5a and Supplementary Table 4). Depression-associated rs1827603 is a brain m⁶A-QTL for the postsynaptic receptor GRM5 (Fig. 5a,b)⁴². Schizophrenia-associated rs7285557 is a brain m⁶A-QTL for a brain-enriched lincRNA,




Fig. 4 | m⁶A QTL versus eQTL comparison. **a**, m⁶A QTLs show a modest but significant enrichment for eQTLs in matched tissues. The enrichment P values were calculated using GARFIELD. $N_{\text{Annot}} = 23,231, 18,663, 43,208, 27,398$ and $44,602$ for Brain Cortex, Brain Frontal Cortex (BA9), Lung, Heart Left Ventricle and Skeletal Muscle (see Methods). The error bars denote the upper and lower bounds for the 95% CI of effect size. The P values were calculated by GARFIELD logistic regression with ‘feature matching’. **b**, Examples of m⁶A QTLs (left) that are also eQTLs (right), with negative (I in **c**) or positive (II in **c**) effects on expression, indicating potential degradation or stabilization effects of m⁶A, respectively. The box shows the 25th–75th percentile; the line shows the median; the whiskers show $1.5 \times \text{IQR}$. **c**, Across all SNPs (dots), only a minority (colored) affect both m⁶A (x axis) and expression (y axis), while most QTLs have independent m⁶A-versus-expression effects (gray points). **d**, Examples of an eQTL with no effect on m⁶A (III in **c**) and an m⁶A QTL with no effect on expression (IV in **c**). The box shows the 25th–75th percentile; the line shows the median; the whiskers show $1.5 \times \text{IQR}$. **e**, A large distance between the lead m⁶A-QTL SNP and the lead eQTL SNP (GTEx v8) for matching target genes indicates distinct mechanisms of action even when both QTL types are identified.

LINC00634, downregulated in schizophrenia⁴³. Anxiety-associated [rs1541627](#) is a brain m⁶A QTL for the synaptic plasticity regulator ANKS1B (ref. 44).

For lung, 62 m⁶A QTLs colocalize with GWAS variants. The peak expiratory flow (PEF)-associated intronic [rs682164](#) SNP is a lung m⁶A QTL for retinoid-responsive CXXC5 that attenuates lung fibrosis in mice⁴⁵. PEF-associated [rs79966207](#) is a lung m⁶A QTL for plexin-B2 (PLXNB2) acting in adult lung bronchial epithelium⁴⁶ (Fig. 5c and Extended Data Fig. 5a). Asthma-associated [rs3194051](#) is a lung m⁶A QTL for immune-related interleukin-7 (IL-7) that contributes to atopic asthma, acting in bronchoalveolar lavage fluid and regulating airway eosinophilia⁴⁷. Forced expiratory volume in 1 s (FEV1)-associated [rs35956171](#) is a lung m⁶A QTL for the alveolar fatty acid oxidation regulator CPT1A, implicated in acute lung injury⁴⁸ (Extended Data Fig. 5a and Supplementary Table 4).

In muscle/heart, 50 m⁶A QTLs colocalize with GWAS variants (Extended Data Fig. 5b). The SNP [rs888298](#), which is associated with coronary artery disease, is a muscle/heart m⁶A QTL targeting the cardiac myocyte mitochondrial oxidation regulator WIP1 involved in signaling and autophagy⁴⁹. The SNP [rs6791834](#), which is associated with heart pulse rate, is a muscle/heart m⁶A QTL for the myocyte microtubule differentiation regulator MAP4 involved in heart development⁵⁰. The SNP [rs56104944](#), which is associated with high blood pressure, is a muscle/heart m⁶A QTL for the heat shock protein HSPA4 involved in cardiac hypertrophy and fibrosis⁵¹ (Extended Data Fig. 5b and Supplementary Table 4).

Beyond GWAS SNPs, m⁶A QTLs helped interpret 3′-UTR and intronic SNPs from ClinVar (Supplementary Table 5). The SNP [rs13069079](#), which is associated with congenital cataracts, is an intronic brain m⁶A QTL for the eye segment morphology regulator FYCO1 (ref. 52; Extended Data Fig. 5c). The SNP [rs605430](#), which is associated with nemaline myopathy and familial restrictive cardiomyopathy, is a 5′-UTR muscle/heart m⁶A QTL for the actin isoform ACTA1, essential for muscle contraction in sarcomeric thin filaments of skeletal muscle⁵³ (Extended Data Fig. 5d).

Tissue-specific m⁶A-QTL enrichments of GWAS variants. We next assessed global tissue-specific m⁶A-QTL enrichments for GWAS variants by heritability partitioning^{54,55} after correcting for eQTLs, and found 27 significantly enriched traits ($P \leq 0.05$; Fig. 5d), indicating that m⁶A QTLs capture residual heritability beyond eQTLs. These results held even after correcting for coding regions, UTRs, promoters and conserved regions (Extended Data Fig. 5e), indicating that they are not simply driven by fortuitous overlaps.

Brain m⁶A QTLs enriched for six traits, including amyotrophic lateral sclerosis, tense mood, schizophrenia and Alzheimer’s disease (AD; Fig. 5d), consistent with roles of m⁶A in brain disorders^{17,21}. Interestingly, AD GWAS variants enrich for GTEx eQTLs⁵⁶ and our m⁶A QTLs in bulk brain samples (composed primarily of neurons), while epigenomic enrichments implicate immune and microglia cells^{57–59}, indicating that both neurons and immune cells may mediate AD genetic effects.

Lung m⁶A QTLs enriched for 19 traits, including both lung-related traits of forced vital capacity, FEV1 and asthma (Fig. 5d), and blood- and immune-related traits, which may reflect immune roles of lung tissue^{60–62}.

Muscle/heart m⁶A QTLs enriched for eight traits, including hypertension, high blood pressure and gout (Fig. 5d), reinforcing reports of m⁶A regulators (FTO, METTL3 and ALKBH5) in cardiovascular diseases⁶³, and suggesting potential interactions between *trans*-regulator effects and *cis*-m⁶A-QTL effects.

Novel m⁶A regulator prediction. As only a small number of m⁶A regulators are known, accounting for only a small fraction of the observed complexity of m⁶A dynamics²², we next used our m⁶A QTLs to help reveal candidate novel m⁶A regulators, by searching for RBPs that preferentially bind m⁶A-QTL regions using data for 171 RBPs across ~34 million target sites from crosslinking and immunoprecipitation (CLIP) with sequencing analyses⁶⁴.

We predicted 69 candidate m⁶A regulators in all three tissues, of which 26 are shared by at least 2 tissues (Bonferroni-adjusted




Fig. 5 | m^{A} -QTLs help interpret GWAS loci. **a, Manhattan plot showing the genomic position (y axis) and m^{A} -QTL P value (x axis) for lead SNPs (points) across traits (colors) showing colocalization of GWAS variants and m^{A} QTLs in brain. **b**, An example of GWAS- m^{A} -QTL colocalization for the depression-associated A-to-G SNP rs1827603, increasing the m^{A} level for *GRM5* in brain, with no effect on *GRM5* expression. The box shows the 25th–75th percentile; the line shows the median; the whiskers show $1.5 \times \text{IQR}$. **c**, An example of GWAS- m^{A} -QTL colocalization for the PEF-related T-to-C SNP rs79966207, increasing the m^{A} level for *PLXNB2* in lung, with no effect on *PLXNB2* expression. The box shows the 25th–75th percentile; the line shows the median; the whiskers show $1.5 \times \text{IQR}$. **d**, GWAS traits (rows) showing significant enrichment (heatmap) for m^{A} QTLs across tissues (columns) by stratified linkage disequilibrium (LD) score regression (S-LDSC) grouped by enriched tissue, and colored by expected tissue of action. The enrichment P value is shown in a color scale; the fold enrichment is shown for those traits with $P < 0.05$. Enrichment P value reported by S-LDSC via z-score calculation, not adjusted for multiple testing. ALS, amyotrophic lateral sclerosis; FVC, forced vital capacity; IBD, inflammatory bowel disease; LDL, low-density lipoprotein.**

$P = 0.05$; Fig. 6a, Extended Data Fig. 6a and Supplementary Table 6), including one m^{A} writer adapter (RBM15), four known m^{A} readers (YTHDF2, YTHDC1, FMR1 and HNRNPC) and four previously proposed candidate m^{A} readers using preferential binding of m^{A} -modified oligonucleotides^{23,24} (CPSF6, NUDT21, TARDBP and PRPF8), consistent with evidence that m^{A} readers can impact the m^{A} level by protecting m^{A} from demethylation by m^{A} erasers^{65,66}.

Our candidates formed a tight protein–protein interaction network with several known m^{A} readers/writers/erasers, suggesting m^{A} cofactor interactions²² reminiscent of pre-transcriptional and splicing regulators. For example, DD3X3 showed experimentally validated interactions with the known m^{A} eraser ALKBH5 (Fig. 6b), with which it was shown to modulate mRNA demethylation⁶⁷. An additional five RBPs (ATXN2, EFTUD2, UPF1, NCBP2 and LARP4; Fig. 6b, dotted circles) showed multiple interactions with both m^{A} writers (purple) and readers (green), indicating that they may function as adapter proteins between them.

We also found several cases where allele-specific RBP binding⁶⁸ overlapped our m^{A} QTLs, including: SRSF1 and PRPF8 for brain m^{A} QTLs; NCBP2, TARDBP, UCHL5 and ZNF622 for lung m^{A} QTLs; and RBM15 for muscle/heart m^{A} QTLs (Supplementary Table 7).

Seven of our 26 candidates were also supported by significant correlations (across individuals, adjusted $q < 0.1$) between RBP expression levels (measured by GTEx RNA-seq) and the m^{A} level of the RBP-bound m^{A} -QTL targets (measured by our eGTEx m^{A} levels; Fig. 6a(ii)). For example, the known m^{A} writer RBM15 and the predicted regulator FIP1L1 (which directly interacts with the known writer ZC3H13) and five additional RBPs (FMR1, NUDT21, STAU1, UCHL5 and ZNF622) were all positively correlated with the m^{A} methylation level of their CLIP-inferred bound targets, consistent with contributions to m^{A} deposition or blocking demethylation (Fig. 6a,c and Extended Data Fig. 6c,d). By contrast, TARDBP, CSTF2T, CPSF7 and GEMIN5 were negatively correlated with the




Fig. 6 | Predicted m⁶A regulators. **a**, Predicted m⁶A regulators (rows) supported by three lines of evidence (check marks): (i) enrichment of RBP regulator-bound sites for m⁶A QTLs in brain (yellow), lung (green) or muscle/heart (purple); (ii) significantly positive (red plus) or negative (green minus) correlation between RBP expression and the methylation level of its m⁶A targets; (iii) experimentally validated protein-protein interaction (PPI) with known m⁶A regulators. **b**, Experimentally determined protein-protein interactions between predicted m⁶A regulators (gray) and known m⁶A regulators (colors) supported by enriched RBP binding in m⁶A QTLs (gray), RBP expression versus m⁶A target level (blue check mark), or direct interaction with known m⁶A regulators (green check mark). Highlighted RBPs (dotted circle) have multiple interactions with both writer adapters and readers, suggesting potential adapter roles. **c**, Examples showing correlation of RBP expression (x axis) versus m⁶A target methylation (y axis) for the predicted regulator STAU1 in brain. Gray shadows denote the 95% confidence region for the regression fit. **d**, Examples showing correlation of RBP expression (x axis) versus m⁶A target methylation (y axis) for the predicted regulator CSTF2T in muscle/heart. Gray shadows denote the 95% confidence region for the regression fit.

m⁶A methylation level of their CLIP-inferred bound targets, consistent with contributions to m⁶A demethylation or blocking deposition (Fig. 6a,d).

Discussion

Our study reports epitranscriptomic interindividual variation in post-transcriptional m⁶A mRNA modifications across multiple primary human tissues, and its integration with genetic variation, expression variation, disease-associated loci and RBP binding, to recognize *cis*- and *trans*-acting drivers of mRNA modifications and their roles in human disease.

We generated transcriptome-wide post-transcriptional modification profiles in humans, spanning 176 experiments across 107 individuals in brain, lung, heart and muscle, filtered to 129 high-quality experiments using enrichment of stop codons and m⁶A consensus motifs and other quality-control metrics. We report 278,000 multiply detected m⁶A sites, greatly expanding the number of known m⁶A-modified loci, and implicating ~43% of all protein-coding genes.

We defined m⁶A QTLs at both a stringent threshold for pinpointing individual m⁶A QTLs and a more inclusive threshold for recognizing functional properties of m⁶A QTLs, as biologically meaningful GWAS enrichments continue well past stringent thresholds to even nominally significant thresholds^{69,70}. We show that m⁶A QTLs at both thresholds are highly reliable, with ~100% directionality consistency between tissues, and significant validation in an independent cohort. Even near-threshold and subthreshold

m⁶A QTLs showed >90% consistency, indicating that even looser thresholds may capture additional reliable and biologically meaningful m⁶A QTLs.

We found that m⁶A QTLs are approximately threefold more tissue specific than eQTLs for matched cohorts, which may stem from tissue-specific co-transcriptional and post-transcriptional m⁶A regulators^{12,71}, and selective pressures against variants with multi-tissue effects, similar to GWAS variants that primarily localize in tissue-specific enhancers rather than tissue-shared promoters. However, the ~470 tissue-shared m⁶A QTLs were >99% consistent in effect directionality between tissues, and even subthreshold m⁶A QTLs were >90% consistent in directionality, indicating that m⁶A *trans*-acting regulators rarely change direction of effect, even though their activity level for different targets may vary across tissues.

We provided several lines of evidence that the observed m⁶A-QTL tissue specificity is biological rather than technical, including: using simulations to assess the m⁶A-QTL calling power, and confirming that our observed data are more tissue specific than expected even at matching MAF, effect size and sample size; partitioning cross-tissue comparisons of m⁶A QTLs across different *P*-value thresholds, and showing that no-effect m⁶A QTLs show near-random directionality, indicating that the lack of m⁶A-QTL effect does not stem from threshold effects; and showing that m⁶A QTLs are substantially more tissue specific than eQTLs, by subsampling GTEx eQTLs to matched individual donors, thus controlling for sample size, discovery power, allele frequency and potential *trans*-acting effects.

We note, however, that mRNA expression measurements are more robust and biologically more stable, while m⁶A may be more variable due to both biological and technical reasons, which may partly account for the observed increased tissue specificity of m⁶A QTLs. Moreover, we expect the fraction of tissue sharing to increase for m⁶A QTLs with increased discovery power at larger sample sizes, as previously observed for eQTLs^{4,8,30}. However, our directionality consistency analysis indicates that the large majority of m⁶A QTLs are tissue specific irrespective of discovery power, as GTEx eQTL directionality shows near-perfect agreement at ~20-fold-reduced sample sizes (for example, from 706 individuals to 38 for muscle eQTLs), thus enabling us to estimate the fraction of true m⁶A-QTL effects even at much lower discovery thresholds.

Our results shed light on ~400 intronic and exonic eQTLs whose mechanism of action was not previously characterized, by showing that they act as tissue-specific m⁶A QTLs, which may mediate their effect on expression levels through mRNA degradation or stability, as previously shown for the YTHDF2 and IGFBP family regulators, respectively. The two directionalities of effect showed uneven proportions among m⁶A QTLs that also impact expression: only one-quarter of increased-m⁶A alleles showed increased expression, indicating roles in mRNA stability, while three-quarters showed decreased expression, indicating roles in mRNA degradation. In the vast majority of cases however, m⁶A QTLs acted through distinct sets of SNPs from eQTLs, thus expanding the set of genetic variants known to affect the molecular phenotypes of protein-coding genes, and helping to shed light on the function of potential disease-associated variants.

Our results also revealed the widespread role of m⁶A QTLs in human disease. At the genome-wide level, we found that m⁶A QTLs were significantly enriched for disease-associated genetic loci and showed compelling tissue-specific enrichments, with psychiatric and neurodegenerative traits enriched in brain m⁶A QTLs, respiratory traits enriched in lung m⁶A QTLs and blood pressure traits enriched in muscle/heart m⁶A QTLs. At the single-locus level, we also found multiple tissue-specific examples of m⁶A QTLs affecting biologically meaningful target genes through colocalization analysis with GWAS SNPs, thus providing new insights and candidate mechanistic hypotheses for GWAS hits that were previously uncharacterized. While experimental validation of these hypotheses through CRISPR–Cas9 genome editing requires tissue systems and animal models of human disease that are not yet developed and will take years to complete for heart, muscle, lung and brain, they can help guide future experiments and pre-clinical studies by expanding the diversity of mechanistic hypotheses underlying human disease genetics to include epitranscriptomic molecular phenotypes, and they demonstrate the broad utility of our results for the field of human genetics and GWAS interpretation.

Our results also revealed 26 RBPs as m⁶A regulator candidates, greatly expanding the set of factors and cofactors currently implicated in m⁶A regulation, which can help guide the systematic dissection of m⁶A regulatory circuitry. All 26 RBPs showed preferential mRNA binding in m⁶A QTLs, 20 also showed protein–protein interactions with known m⁶A regulators, 7 also showed allele-specific RBP binding overlapping our m⁶A QTLs, and 11 also showed significant expression correlation with the m⁶A levels of their targets. Five of the putative novel regulators (UPF1, NCBP2, LARP4, ATXN2 and EFTUD2) were particularly intriguing, as they showed multiple interactions with both m⁶A writers and readers, indicating possible functions as adapter proteins between them. The tight network of our predicted m⁶A regulators and known m⁶A regulators suggests extensive cooperation of multiple cofactors guiding m⁶A methylation, as in current pre-transcriptional and splicing regulatory models.

Our study has several limitations. First, our study is still confined to a relatively small number of tissues and a small number of individuals, due to the difficulty and sample requirements of profiling

m⁶A in primary human tissues. Thus, our study identified only the strongest m⁶A QTLs, and we expect future studies to reveal many more m⁶A QTLs, which we estimate will show a modestly higher overlap between tissues, although still substantially lower than for eQTLs. Second, our antibody-based m⁶A profiling does not provide single-nucleotide resolution and is not fully quantitative. We expect future studies to overcome these limitations, although existing methods that rely on restriction enzymes are confined to specific sequence contexts^{72–76} and can detect only a subset of m⁶A sites. Third, similar to GTEx, our study sampled m⁶A levels only at bulk-tissue resolution, without distinguishing between different cell types within them. We expect that single-cell profiling of m⁶A modifications, when such technologies become available, will help reveal the specific cell types where our m⁶A QTLs act, and to also reveal additional single-cell m⁶A QTLs that are not captured at the bulk level. Last, we focused on polyadenylated protein-coding mRNAs, while m⁶A methylation can affect chromatin-associated RNAs and other non-coding RNAs, whose interindividual m⁶A variation may be highly informative in gene regulation and disease studies.

Overall, our datasets and analyses provide a foundation to bridge genetic variation with epitranscriptomic regulation and human disease in human primary tissues. This is particularly important as m⁶A is increasingly recognized to play important roles in human disease, and many disease-associated variants remain uninterpreted. The resulting m⁶A QTLs, target genes, upstream regulators and biological insights can help further our understanding of epitranscriptomic gene-regulatory control, and help pave the way for new therapeutic targets in human disease.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41588-021-00890-3>.

Received: 5 September 2020; Accepted: 18 May 2021;

Published online: 1 July 2021

References

1. Ward, L. D. & Kellis, M. Interpreting noncoding genetic variation in complex traits and human disease. *Nat. Biotechnol.* **30**, 1095–1106 (2012).
2. Tak, Y. G. & Farnham, P. J. Making sense of GWAS: using epigenomics and genome engineering to understand the functional relevance of SNPs in non-coding regions of the human genome. *Epigenetics Chromatin* **8**, 57 (2015).
3. Grundberg, E. et al. Mapping *cis*- and *trans*-regulatory effects across multiple tissues in twins. *Nat. Genet.* **44**, 1084–1089 (2012).
4. Lonsdale, J. et al. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–585 (2013).
5. Sun, W. et al. Histone acetylome-wide association study of autism spectrum disorder. *Cell* **167**, 1385–1397 (2016).
6. Wainberg, M. et al. Opportunities and challenges for transcriptome-wide association studies. *Nat. Genet.* **51**, 592–599 (2019).
7. Huan, T. et al. Genome-wide identification of DNA methylation QTLs in whole blood highlights pathways for cardiovascular disease. *Nat. Commun.* **10**, 4267 (2019).
8. GTEx Consortium. Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).
9. Wu, L. et al. Variation and genetic control of protein abundance in humans. *Nature* **499**, 79–82 (2013).
10. Li, Y. I. et al. RNA splicing is a primary link between genetic variation and disease. *Science* **352**, 600–604 (2016).
11. Park, E. et al. Population and allelic variation of A-to-I RNA editing in human transcriptomes. *Genome Biol.* **18**, 143 (2017).
12. Roundtree, I. A., Evans, M. E., Pan, T. & He, C. Dynamic RNA modifications in gene expression regulation. *Cell* **169**, 1187–1200 (2017).
13. Frye, M., Jaffrey, S. R., Pan, T., Rechavi, G. & Suzuki, T. RNA modifications: what have we learned and where are we headed? *Nat. Rev. Genet.* **17**, 365–372 (2016).

14. Zaccara, S., Ries, R. J. & Jaffrey, S. R. Reading, writing and erasing mRNA methylation. *Nat. Rev. Mol. Cell Biol.* **20**, 608–624 (2019).
15. Zhao, B. S., Roundtree, I. A. & He, C. Post-transcriptional gene regulation by mRNA modifications. *Nat. Rev. Mol. Cell Biol.* **18**, 31–42 (2017).
16. Barbieri, I. & Kouzarides, T. Role of RNA modifications in cancer. *Nat. Rev. Cancer* <https://doi.org/10.1038/s41568-020-0253-2> (2020).
17. Yao, B. et al. Epigenetic mechanisms in neurogenesis. *Nat. Rev. Neurosci.* **17**, 537–549 (2016).
18. Li, X., Xiong, X. & Yi, C. Epitranscriptome sequencing technologies: decoding RNA modifications. *Nat. Methods* **14**, 23–31 (2016).
19. Frye, M., Harada, B. T., Behm, M. & He, C. RNA modifications modulate gene expression during development. *Science* **361**, 1346–1349 (2018).
20. Huang, H., Weng, H. & Chen, J. m⁶A modification in coding and non-coding RNAs: roles and therapeutic implications in cancer. *Cancer Cell* **37**, 270–288 (2020).
21. Livneh, I., Moshitch-Moshkovitz, S., Amariglio, N., Rechavi, G. & Dominissini, D. The m⁶A epitranscriptome: transcriptome plasticity in brain development and function. *Nat. Rev. Neurosci.* **21**, 36–51 (2020).
22. Shi, H., Wei, J. & He, C. Where, when, and how: context-dependent functions of RNA methylation writers, readers, and erasers. *Mol. Cell* **74**, 640–650 (2019).
23. Arguello, A. E., DeLiberto, A. N. & Kleiner, R. E. RNA chemical proteomics reveals the N⁶-methyladenosine (m⁶A)-regulated protein–RNA interactome. *J. Am. Chem. Soc.* **139**, 17249–17252 (2017).
24. Edupuganti, R. R. et al. N⁶-methyladenosine (m⁶A) recruits and repels proteins to regulate mRNA homeostasis. *Nat. Struct. Mol. Biol.* **24**, 870–878 (2017).
25. Patil, D. P., Pickering, B. F. & Jaffrey, S. R. Reading m⁶A in the transcriptome: m⁶A-binding proteins. *Trends Cell Biol.* **28**, 113–127 (2018).
26. Liu, J. et al. Landscape and regulation of m⁶A and m⁶Am methylome across human and mouse tissues. *Mol. Cell* **77**, 426–440 (2020).
27. Xiao, S. et al. The RNA N⁶-methyladenosine modification landscape of human fetal tissues. *Nat. Cell Biol.* **21**, 651–661 (2019).
28. Zhang, Z. et al. Genetic analyses support the contribution of mRNA N⁶-methyladenosine (m⁶A) modification to human disease heritability. *Nat. Genet.* **52**, 939–949 (2020).
29. eGTEx Project. Enhancing GTEx by bridging the gaps between genotype, gene expression, and disease. *Nat. Genet.* **49**, 1664–1670 (2017).
30. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318–1330 (2020).
31. Dominissini, D. et al. Topology of the human and mouse m⁶A RNA methylomes revealed by m⁶A-seq. *Nature* **485**, 201–206 (2012).
32. Schwartz, S. et al. High-resolution mapping reveals a conserved, widespread, dynamic mRNA methylation program in yeast meiosis. *Cell* **155**, 1409–1421 (2013).
33. Xuan, J.-J. et al. RMBase v2.0: deciphering the map of RNA modifications from epitranscriptome sequencing data. *Nucleic Acids Res.* **46**, D327–D334 (2018).
34. Meyer, K. D. et al. Comprehensive analysis of mRNA methylation reveals enrichment in 3′ UTRs and near stop codons. *Cell* **149**, 1635–1646 (2012).
35. Lin, S., Choe, J., Du, P., Triboulet, R. & Gregory, R. I. The m⁶A methyltransferase METTL3 promotes translation in human cancer cells. *Mol. Cell* **62**, 335–345 (2016).
36. Visvanathan, A. et al. Essential role of METTL3-mediated m⁶A modification in glioma stem-like cells maintenance and radioresistance. *Oncogene* **37**, 522–533 (2018).
37. Stegle, O., Parts, L., Durbin, R. & Winn, J. A Bayesian framework to account for complex non-genetic factors in gene expression levels greatly increases power in eQTL studies. *PLoS Comput. Biol.* **6**, e1000770 (2010).
38. Lappalainen, T. et al. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature* **501**, 506–511 (2013).
39. Huang, H. et al. Recognition of RNA N⁶-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. *Nat. Cell Biol.* **20**, 285–295 (2018).
40. Wang, X. et al. N⁶-methyladenosine-dependent regulation of messenger RNA stability. *Nature* **505**, 117–120 (2014).
41. Amaral, A. C. et al. Prenatal protein malnutrition decreases *KCNJ3* and 2DG activity in rat prefrontal cortex. *Neuroscience* **286**, 79–86 (2015).
42. Doré, A. S. et al. Structure of class C GPCR metabotropic glutamate receptor 5 transmembrane domain. *Nature* **511**, 557–562 (2014).
43. Gandal, M. J. et al. Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* **362**, eaat8127 (2018).
44. Tindi, J. O. et al. *ANKK1B* gene product AIDA-1 controls hippocampal synaptic transmission by regulating GluN2B subunit localization. *J. Neurosci.* **35**, 8986–8996 (2015).
45. Cheng, W., Wang, F., Feng, A., Li, X. & Yu, W. CXXC5 attenuates pulmonary fibrosis in a bleomycin-induced mouse model and MLFs by suppression of the CD40/CD40L pathway. *Biomed Res. Int.* **2020**, 7840652 (2020).
46. Zielonka, M., Xia, J., Friedel, R. H., Offermanns, S. & Worzfeld, T. A systematic expression analysis implicates Plexin-B2 and its ligand Sema4C in the regulation of the vascular and endocrine system. *Exp. Cell Res.* **316**, 2477–2486 (2010).
47. Kelly, E. A. B. et al. Potential contribution of IL-7 to allergen-induced eosinophilic airway inflammation in asthma. *J. Immunol.* **182**, 1404–1410 (2009).
48. Cui, H. et al. Impairment of fatty acid oxidation in alveolar epithelial cells mediates acute lung injury. *Am. J. Respir. Cell Mol. Biol.* **60**, 167–178 (2019).
49. Tzimas, C. et al. WIPI1 is a conserved mediator of right ventricular failure. *JCI Insight* **5**, e122989 (2019).
50. Webster, D. R. & Bratcher, J. M. Developmental regulation of cardiac MAP4 protein expression. *Cell Motil. Cytoskeleton* **63**, 512–522 (2006).
51. Mohamed, B. A. et al. Targeted disruption of *Hspa4* gene leads to cardiac hypertrophy and fibrosis. *J. Mol. Cell. Cardiol.* **53**, 459–468 (2012).
52. Hernandez-Zimbron, L. F. et al. Molecular age-related changes in the anterior segment of the eye. *J. Ophthalmol.* **2017**, 1295132 (2017).
53. Laing, N. G. et al. Mutations and polymorphisms of the skeletal muscle α -actin gene (*ACTA1*). *Hum. Mutat.* **30**, 1267–1277 (2009).
54. Finucane, H. K. et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
55. Andersson, R. et al. An atlas of active enhancers across human cell types and tissues. *Nature* **507**, 455–461 (2014).
56. Gamazon, E. R. et al. Using an atlas of gene regulation across 44 human tissues to inform complex disease- and trait-associated variation. *Nat. Genet.* **50**, 956–967 (2018).
57. Roadmap Epigenomics Consortium et al. Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015).
58. Gjonneska, E. et al. Conserved epigenomic signals in mice and humans reveal immune basis of Alzheimer’s disease. *Nature* **518**, 365–369 (2015).
59. Corces, M. R. et al. Single-cell epigenomic analyses implicate candidate causal variants at inherited risk loci for Alzheimer’s and Parkinson’s diseases. *Nat. Genet.* **52**, 1158–1168 (2020).
60. Ardain, A., Marakalala, M. J. & Leslie, A. Tissue-resident innate immunity in the lung. *Immunology* **159**, 245–256 (2020).
61. Whitsett, J. A. & Alenghat, T. Respiratory epithelial cells orchestrate pulmonary innate immunity. *Nat. Immunol.* **16**, 27–35 (2015).
62. Chiu, C. & Openshaw, P. J. Antiviral B cell and T cell immunity in the lungs. *Nat. Immunol.* **16**, 18–26 (2015).
63. Zheng, N., Su, J., Hu, H., Wang, J. & Chen, X. Research progress of N⁶-methyladenosine in the cardiovascular system. *Med. Sci. Monit.* **26**, e921742 (2020).
64. Zhu, Y. et al. POSTAR2: deciphering the post-transcriptional regulatory logics. *Nucleic Acids Res.* **47**, D203–D211 (2019).
65. Zhou, J. et al. Dynamic m⁶A mRNA methylation directs translational control of heat shock response. *Nature* **526**, 591–594 (2015).
66. Yang, Y., Hsu, P. J., Chen, Y.-S. & Yang, Y.-G. Dynamic transcriptomic m⁶A decoration: writers, erasers, readers and functions in RNA metabolism. *Cell Res.* **28**, 616–624 (2018).
67. Shah, A. et al. The DEAD-box RNA helicase DDX3 interacts with m⁶A RNA demethylase ALKBH5. *Stem Cells Int.* **2017**, 8596135 (2017).
68. Yang, E.-W. et al. Allele-specific binding of RNA-binding proteins reveals functional genetic variants in the RNA. *Nat. Commun.* **10**, 1338 (2019).
69. Wang, X. et al. Discovery and validation of sub-threshold genome-wide association study loci using epigenomic signatures. *eLife* **5**, e10557 (2016).
70. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An expanded view of complex traits: from polygenic to omnigenic. *Cell* **169**, 1177–1186 (2017).
71. Zhou, K. I. et al. Regulation of co-transcriptional pre-mRNA splicing by m⁶A through the low-complexity protein hnRNPG. *Mol. Cell* **76**, 70–81 (2019).
72. Garcia-Campos, M. A. et al. Deciphering the ‘m⁶A code’ via antibody-independent quantitative profiling. *Cell* **178**, 731–747 (2019).
73. Wang, Y., Xiao, Y., Dong, S., Yu, Q. & Jia, G. Antibody-free enzyme-assisted chemical approach for detection of N⁶-methyladenosine. *Nat. Chem. Biol.* <https://doi.org/10.1038/s41589-020-0525-x> (2020).
74. Zhang, Z. et al. Single-base mapping of m⁶A by an antibody-independent method. *Sci. Adv.* **5**, eaax0250 (2019).
75. Shu, X. et al. A metabolic labeling method detects m⁶A transcriptome-wide at single base resolution. *Nat. Chem. Biol.* <https://doi.org/10.1038/s41589-020-0526-9> (2020).
76. Liu, H. et al. Accurate detection of m⁶A RNA modifications in native RNA sequences. *Nat. Commun.* **10**, 4079 (2019).

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature America, Inc. 2021

GTEx Consortium

Kristin G. Ardlie² and François Aguet²

Methods

m⁶A meRIP-seq across human tissues. Sample collection by the GTEx Consortium, with donor enrollment and consent, histopathological review procedures, biospecimen procurement methods and fixation, and informed consent approval, was as previously described⁷⁷. m⁶A profiling was performed across four eGTEx tissues, including Brain Cortex (*n* = 53), Lung (*n* = 32), Skeletal Muscle (*n* = 32) and Heart Left Ventricle (*n* = 12), using an optimized version of meRIP-seq. These four tissue types were chosen as they are highly disease relevant and are representative for different germ layers (brain: ectoderm; muscle/heart: mesoderm; lung: endoderm). To minimize potential confounding factors introduced by experimental differences, the samples from each tissue were balanced across three experimental batches (Supplementary Table 1). We adopted the protocols from Batista et al.⁷⁸, adapted to enable 400-fold less starting material available for the eGTEx samples. We calibrated the m⁶A-Dynabeads complex by coupling Synaptic System anti-m⁶A to magnetic beads at a ratio of 5 μg of antibody to 1 mg of Dynabeads, and adjusting the reaction conditions and the volume of the m⁶A-Dynabeads complex to 25 μl in a total volume of 50 μl for immunoprecipitation (IP). The new wash steps were set at 100 μl volume, and the new elution steps were established at 4 consecutive elution steps each using 50 μl of elution buffer at 52 °C for 5 min under gentle rotation. The combined 240 μl elution was then precipitated by the addition of 400 mM NaCl, 2 μl of ultrapure glycogen at 20 mg ml⁻¹ and 2.5 times the volume of 200-proof ethanol at -20 °C for 1 h or overnight. The RNA was precipitated at 4 °C for 25 min by centrifugation at 13,000g. The pellet was then washed twice in 70% ethanol, dried at 25 °C for 15 min, and resuspended in 7 μl of ultrapure H₂O before library construction. We used the SMARTer Stranded RNA-Seq Kit from Clontech/Takada, which is optimized to work with 100 pg of starting material. The libraries were sent for 2 × 45-base-pair paired-end sequencing.

m⁶A site identification and quantification. The RNA-seq (without immunoprecipitation) data were taken from the GTEx portal (<https://gtexportal.org/home/datasets>) as input for m⁶A peak calling and quantification. Reads from the RNA-seq and meRIP-seq were first aligned against tRNA (downloaded from the University of California, Santa Cruz Table Browser) and rRNA (downloaded from the National Center for Biotechnology Information Nucleotide database) using bowtie2 (v2.3.4.3)⁷⁹, with the unmapped reads kept for further analysis. The remaining reads were mapped to the hg38 human genome using hisat2 (v2.1.0)⁸⁰, with GENCODE (v26; downloaded from <https://gtexportal.org/home/datasets>) as annotation, and with the following parameters: -k 1 --no-discordant. Peak calling was carried out using MACS2 (version 2.1.1), with the parameters of --nomodel and --extsize 100. Peak processing was performed using bedtools (v2.28.0). The high-confidence peaks were selected by requiring *q* value ≤ 10⁻⁵ and fold change ≥ 3. The raw methylation level for each m⁶A site was quantified as²³:

$$\log \left[\frac{\text{Reads}_{\text{Peak,meRIP}} / \text{Reads}_{\text{Peak,RNA-seq}}}{\text{Reads}_{\text{Total,meRIP}} / \text{Reads}_{\text{Total,RNA-seq}}} \right]$$

The m⁶A level matrix across samples and peaks was further normalized before m⁶A-QTL calling (see the section below entitled m⁶A normalization).

m⁶A site quality-control metrics. To evaluate whether the m⁶A immunoprecipitation was successful, we used deepTools (v3.0.2) to examine the read density of meRIP-seq surrounding the GGACH motif, compared to other control sequence motifs, including the reverse complement sequence (DGTC), reverse sequence (HCAGG) and complement sequence (CCTGD). We then evaluated the quality of the identified m⁶A peaks using the two well-recognized m⁶A metrics: m⁶A consensus sequence and distribution along mRNA. For the distribution evaluation, we first converted the genomic m⁶A positions to transcriptomic coordinates, and then calculated the density of m⁶A along the mRNA structure. For each individual sample, we performed motif analysis using AME (v5.0.3) from the MEME tool by examining the enrichment of the m⁶A motif (GGACH) over the randomly shuffled sequences with the nucleotide content preserved. The enrichment *P* value of the motif was calculated by AME based on Fisher's exact test, and multiple-testing correction was performed using Bonferroni adjustment.

Tissue specificity of m⁶A sites. We performed hierarchical clustering with the global sequencing signals by segmenting the genome into 200-kb bins. For the individual m⁶A sites, we defined an m⁶A site as tissue-specific m⁶A if it occurred in only one of the four tissues. We then further categorized the tissue-specific m⁶A into two groups: expression-specificity-induced tissue-specific m⁶A and epitranscriptome-layer tissue-specific m⁶A, depending on whether the tissue specificity mainly comes from the gene expression level or not. A tissue-specific m⁶A is considered as an expression-specificity-induced tissue-specific m⁶A when the expression of the host gene in the tissue is ten times higher than in any of the other tissues, whereas the rest were defined as epitranscriptome-layer tissue-specific m⁶A.

m⁶A and SNP pre-filtration. For each tissue, we filtered out the variants with fewer than 8 minor allele individuals (minor allele homozygotes + heterozygotes)

to avoid false signals caused by a small number of individuals and to lower the multi-test correction burden. Additionally, we envision that m⁶A regulation may take place: at the DNA level, where a SNP in DNA may affect m⁶A given that the installation process occurs co-transcriptionally; at the RNA level, where a SNP in a transcript can modulate the m⁶A level in *cis* in the same RNA molecule in a post-transcriptional fashion. Therefore, to ensure both the sensitivity and the biological rationality, we specified the m⁶A-QTL searching regions to be the promoter plus the gene body for each m⁶A in a gene.

For m⁶A-QTL calling, we selected the high-confidence m⁶A sites by requiring either that the m⁶A peak was previously curated³³ and meanwhile was captured in a tissue for no fewer than two individuals; or that the m⁶A peak was captured across 20% of the samples in a tissue.

m⁶A normalization. The m⁶A signal matrix of one tissue was further processed and normalized before QTL identification: m⁶A sites with more than 20% of the values missing (caused by undetectable RNA expression) across individuals were excluded; the IP efficiency differences between samples and the GC content across peaks were corrected following well-established procedures²⁸; the IP- and GC-corrected m⁶A matrix was further standardized by subtracting the mean plus dividing by the s.d. for each peak across all of the samples in that tissue; last, the matrix was quantile-normalized across all of the peaks. The resulting m⁶A matrix was used for latent factor identification and m⁶A-QTL calling.

Latent factor identification and removal. To regress out the unwanted variance stemming from non-genetic factors, we employed PEER³⁷ (R package 1.0) to identify the latent covariates for the samples in each tissue. We chose the PEER factors for QTL calling based on: maximizing the number of g-m⁶A sites identified (Fig. 2b); and the correlation between the PEER factors (Fig. 2a and Extended Data Fig. 2a,b). We included the top seven, two and five PEER factors for brain, lung and muscle/heart, respectively, plus experimental batch and the top three genotype PCs as covariates for m⁶A-QTL calling.

m⁶A-QTL identification. We used FastQTL (ref.⁸¹; v2.0) for m⁶A-QTL identification with the PEER factors included as covariates as described above. The searching window was set to the gene body plus the promoter (1 kb upstream of the transcription start site (TSS)). To identify g-m⁶A sites, the nominal *P* value of the lead SNP of each m⁶A peak was corrected to get an empirical *P* value that accounts for the multiple variants tested. The empirical *P* value was extrapolated based on beta distribution fitting to the permutation results by FastQTL (parameter --permutate, 1,000). We applied a threshold of empirical *P* value < 0.005 on the lead variant to identify g-m⁶A sites, and then for each g-m⁶A we applied the nominal *P* value threshold that corresponds to the empirical *P* value of 0.005 for each locus to identify m⁶A QTLs (Supplementary Table 2). Given the larger sample size in brain, we provided higher-confidence sets of g-m⁶A sites and m⁶A QTLs by further correcting the empirical *P* value with the Benjamini-Hochberg procedure and applying a threshold of FDR < 0.2 (Supplementary Table 3).

m⁶A-QTL validation. To validate the m⁶A QTLs we identified in an unbiased way, we utilized an independent dataset published by another study²⁶, from which we obtained the m⁶A profiles of heart, muscle and lung, each with two adult post-mortem individuals. Although brain samples are available from this study, the region they used was from cerebrum, which is much larger and more complex than the cortex region used by our m⁶A-QTL analysis. Given the complexity of the human brain, we did not include the brain samples in the validation. To perform the validation, we first grouped the m⁶A QTLs from our study based on the direction of effect size, where '+' represents the m⁶A level increases from major to minor alleles, and '-' represents the opposite, together with a 'control' group of SNPs that are not m⁶A QTLs. We then used samtools (v1.9) to call SNPs with RNA-seq data of each individual from the validation study. The SNPs with different genotypes in the two individuals were kept for further usage. For each SNP, the individual with more minor alleles was regarded as 'minor' while the other one was regarded as 'major'. The m⁶A fold change of the 'minor' individual over the 'major' one was calculated for each variant in the validation cohort, and compared across the '-', '+' and 'control' groups defined in this study by a Wilcoxon rank-sum test.

Simulation analysis for m⁶A-QTL calling power evaluation. We carried out simulations to assess the power of m⁶A-QTL calling, and thus to assess the extent of m⁶A-QTL tissue specificity driven by the current sample sizes, following the simulation strategy from a published eQTL power analysis study⁸². Specifically, we used the effect size, MAF and sample size parameters observed from the lead m⁶A QTLs we identified, and performed the same number of simulations as the number of g-m⁶A sites identified in each tissue following a simple linear regression: m⁶A = βg + ε, where ε ~ N(0, 1) and g denotes genotypes, assuming that each g-m⁶A site is regulated by only one SNP. We also simulated the same number of controls with the effect size set to 0. For m⁶A QTLs identified in each tissue, we simulated a corresponding set of m⁶A sites with the sample sizes of other tissues. m⁶A QTLs were mapped with linear regression in the simulation dataset. The m⁶A-QTL overlap rate and directionality consistency were calculated by comparing

the simulated dataset with the real dataset from which the m⁶A-QTL parameters were used for simulation.

Enrichment testing and overlap analysis between m⁶A QTLs and eQTLs. The enrichment analysis of m⁶A QTLs versus eQTLs was carried out using GARFIELD (ref.⁸³; v2). The latest release of eQTL data was used (v8), available in the GTEx portal (<https://gtexportal.org/home/>). The enrichment of m⁶A QTLs versus eQTLs was quantified with odds ratios, and the significance was calculated using a generalized linear model, with MAF, distance to TSS and number of LD proxies ($r^2 > 0.8$) accounted for (see the GARFIELD manual for details). To test whether the overlap between eGenes and gmGenes is more than expected by chance (Fig. 4e), we randomly sampled the same number of eGenes and gmGenes from the expressed gene pool in our data ($n = 26,359$) 10,000 times; the P value was calculated as the percentage of cases where the overlap between the randomly sampled genes was greater than the observed overlap.

To enable a more accurate comparison between the effect of m⁶A QTLs on m⁶A versus the effect of eQTLs on the corresponding gene expression, we re-performed eQTL analysis in each tissue using the same cohort as used in m⁶A-QTL calling. As in the m⁶A-QTL analysis, we first used PEER to capture the unwanted variance and included the first 6, 3 and 4 PEER factors for brain, lung and muscle/heart as covariates during eQTL calling. The slopes of eQTLs and m⁶A QTLs were then compared. Correlation of the slopes between the overlapped m⁶A QTLs and eQTLs (separated by effect directionality) was calculated using a two-sided Pearson correlation test; the P value was calculated with linear regression.

GWAS heritability partition of m⁶A QTLs. Enrichment analysis of m⁶A QTLs in GWAS was performed using S-LDSC (v1.0.1) based on the tutorial in ref.⁵⁴. The enrichment was calculated as the proportion of heritability over the proportion of SNPs, and then the standard error was estimated and used for P -value calculation. We used two baseline models: baseline v1.2 (provided by LDSC) plus eQTLs from the same tissue (Extended Data Fig. 5e); baseline v1.2 with annotations related to the gene body excluded (coding region, UTRs, promoter and conserved region), and with the eQTLs from the same tissue included (Fig. 5d). We used two sets of m⁶A QTLs with different thresholds: $P < 10^{-3}$ and $P < 10^{-4}$. For brain m⁶A-QTL heritability analysis, we further included the high-confidence m⁶A QTLs that were identified with two rounds of multiple-testing correction (labeled as '2StepAdj.' in the figure). The GWAS summary statistics files were downloaded from the Alkes group (https://data.broadinstitute.org/alkesgroup/LDSCORE/independent_sumstats/) and UK Biobank; the respiratory traits were downloaded from the GWAS Atlas. Disease-relevant traits were selected.

Colocalization analysis between m⁶A QTLs and GWAS loci. Colocalization analysis between m⁶A QTLs and GWAS variants was carried out using a Bayesian framework for colocalization (Coloc⁸⁴). GWAS loci (filtered by $P < 10^{-4}$) that overlap with g-m⁶A sites were used for colocalization tests. The ratio between PP3 (posterior probability of both traits being associated with different causal variants) and PP4 (posterior probability of both traits being associated with the same causal variant) was calculated, and those loci with PP3/PP4 < 0.05 and PP4 > 0.1 were selected as the GWAS–m⁶A-QTL colocalized events.

Enrichment analysis of m⁶A QTLs in RBP binding. To predict potential indirect m⁶A regulators that function through binding to m⁶A QTLs, we carried out enrichment analysis between m⁶A QTLs and RBP-binding sites. We downloaded the binding sites of 171 RBPs curated by the POSTAR2 database⁶⁴, and then performed enrichment analysis against m⁶A QTLs using GARFIELD⁸³. The enrichment of m⁶A QTLs in RBP-binding sites was quantified as an odds ratio, and the significance was calculated using the generalized linear model, with MAF, distance to TSS and number of LD proxies ($r^2 > 0.8$) accounted for. RBPs with fewer than 100 baseline SNPs overlapping binding sites (NAnnot < 100) were excluded to ensure detection reliability. RBPs showing a Bonferroni-adjusted P value < 0.05 were selected as potential m⁶A regulators.

To further prioritize the enriched RBPs, we carried out a RBP expression versus m⁶A level correlation analysis. For each enriched RBP, we first calculated an 'aggregated' m⁶A score for each individual, where the methylation levels of the m⁶A sites whose corresponding m⁶A QTLs are within the RBP-binding sites were aggregated, and then performed regression between the aggregated m⁶A level and the gene expression level of the RBP across each individual. P values were calculated based on linear regression, and multiple-testing correction was performed by Benjamini–Hochberg (BH) correction. An FDR-adjusted P value < 0.1 was used as the threshold.

Protein–protein interaction analysis. We examined the protein–protein interaction network composed of the enriched RBPs and the known m⁶A regulators (writer/reader/eraser) using the STRINGdb R package (v1.24.0)⁸⁵. Only interactions with 'Experiments' as the source were considered.

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

All eGTEx protected data, including m⁶A sequencing reads and matched RNA-seq data, are available on dbGaP with the accession number [phs000424.v8.p2](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=phs000424.v8.p2). Additionally, the data can be accessed via AnVIL with authentication: https://anvil.terra.bio/#workspaces/anvil-datastorage/AnVIL_GTEx_V8_hg38. As the raw sequencing data with genetic information are protected, application and authentication are needed before accessing the data. All non-protected data for m⁶A can be visualized via the GTEx portal (<https://www.gtexportal.org>) as part of eGTEx v8. The m⁶A QTLs identified in each tissue can be downloaded from Supplementary Tables 1–7. The eQTL datasets are from GTEx v8, which can be accessed at <https://gtexportal.org/home/datasets>. The LCL m⁶A-QTL datasets from Zhang et al.²⁸ can be downloaded from <https://doi.org/10.5281/zenodo.3870952>. The previously curated m⁶A sites can be downloaded from RMBase (<http://rna.sysu.edu.cn/rmbase/>). The RNA-binding sites can be downloaded from <http://lulab.life.tsinghua.edu.cn/postar/>.

Code availability

Code for m⁶A data processing, m⁶A-QTL calling and relevant functional analyses and additional information can be found at <http://compbio.mit.edu/m6AQTLs/> and also on Zenodo at <https://doi.org/10.5281/zenodo.4764136>.

References

- Carithers, L. J. et al. A novel approach to high-quality postmortem tissue procurement: the GTEx Project. *Biopreserv. Biobank.* **13**, 311–319 (2015).
- Batista, P. J. et al. m⁶A RNA modification controls cell fate transition in mammalian embryonic stem cells. *Cell Stem Cell* **15**, 707–719 (2014).
- Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).
- Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* **12**, 357–360 (2015).
- Ongen, H., Buil, A., Brown, A. A., Dermizakis, E. T. & Delaneau, O. Fast and efficient QTL mapper for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479–1485 (2016).
- Huang, Q. Q., Ritchie, S. C., Brozynska, M. & Inouye, M. Power, false discovery rate and Winner's Curse in eQTL studies. *Nucleic Acids Res.* **46**, e133 (2018).
- Iotchkova, V. et al. GARFIELD classifies disease-relevant genomic features through integration of functional annotations with association signals. *Nat. Genet.* **51**, 343–353 (2019).
- Giambartolomei, C. et al. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).
- Szklarczyk, D. et al. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**, D607–D613 (2019).

Acknowledgements

We thank C. Boix for helping with figure visualization, analysis suggestion and discussion, and K. Kang and other Kellis laboratory members for discussions and suggestions. We thank the GTEx Consortium for providing the samples for m⁶A profiling. We thank K. G. Ardlie and F. Aguet for helping with GTEx data deposition and providing helpful suggestions. This work was supported by the NIH grants NIH HG007610, HG008155, HG009446, MH109978, AG054012, AG058002, AG062377, NS110453, NS115064, AG067151, AG062335 and MH119509 to M.K., and CA232115 and CA233671 to R.I.G.

Author contributions

This study was designed by X.X., L.H. and M.K., and directed and coordinated by M.K. B.M. performed the m⁶A profiling; X.X. and L.H. performed the bioinformatic analysis with help from Y.P. and R.I.G. and under the supervision of M.K. All authors participated in the discussion of the project. X.X., L.H. and M.K. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Additional information


Extended data is available for this paper at <https://doi.org/10.1038/s41588-021-00890-3>.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41588-021-00890-3>.

Correspondence and requests for materials should be addressed to M.K.


Peer review information *Nature Genetics* thanks the anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.




Extended Data Fig. 1 | See next page for caption.

Extended Data Fig. 1 | m⁶A landscape across tissues. **a.** Positional enrichment for aggregated meRIP-seq signal surrounding the known m⁶A motif (GGACH) vs. three control sequences (colors). **b.** m⁶A peak positional density (y-axis) along the gene structure (x-axis) in brain, lung, heart and muscle. **c.** Cumulative distribution (y-axis) of GGACH motif enrichment (-log₁₀q-value, x-axis) across samples (teal) vs. shuffled controls (salmon). **d.** Peak Count per sample. **e.** Peaks shared by fewer individuals (left) are more likely previously-undetected (red). **f.** Similar to b for only previously-unreported m⁶A peaks found in >=2 individuals in our study. **g.** Pearson correlation (heatmap) and hierarchical clustering (tree) of mRNA-seq profiles across the 129 GTEx samples (rows/columns) that match the individuals profiled here shows tissue-specific clustering (colors) and co-clustering of heart and muscle, as shown for m⁶A profiles in Fig. 1b. **h.** KEGG pathway enrichments for the genes harboring tissue-specific m⁶A.




Extended Data Fig. 2 | Identification of genetically-driven m⁶A. **a.** Pearson correlation (color) for PEER factors shows saturation (correlated factors) after first 2 factors (red box) in lung (top) and after first 5 factors (red box) in muscle/heart (bottom). **b.** Pearson correlation between PEER factors (columns) and donor phenotypic measurements (rows) helps interpret factors in lung and muscle/heart. **c,d.** Manhattan plot of m⁶A-QTLs in lung (c) and muscle/heart (d), and m⁶A-QTL examples. Boxes=25%-75% percentile; line=median; whiskers=1.5 IQR; p-values=FastQTL linear regressions.




Extended Data Fig. 3 | See next page for caption.

Extended Data Fig. 3 | Tissue specificity of m⁶A-QTL. **a.** Summary of shared/specific g-m⁶As and m⁶A across tissues. **b.** KEGG pathway enrichments for tissue-specific gmGenes. **c.** Tissue-intersections of the eQTLs identified from the same samples for m⁶A-QTL calling. **d,e.** Correlation between p-value (x-axis) and eQTL effect (y-axis) between tissues, with positive- and negative-effect eQTLs separated for full GTEx-V8 (d), and subsampled to individuals used here (e);. **f.** Nominal p-values (y-axis) of simulated m⁶A-QTLs (Positive), and simulated NULL m⁶A-QTLs controls without QTL effects (Control). Boxes=25%-75% percentile; line=median; whiskers=1.5 IQR. **g.** m⁶A-QTL overlaps between tissues in simulated data show much higher tissue-sharing (teal curve) than in observed real data (peach curve). **h.** Effect sizes directionality between m⁶A-QTL from real tissue data and simulated data are almost 100% consistent. **i.** Effect directionality consistency when m⁶A-QTLs were identified with 50, 38 and 30 samples. **j.** Gene expression distribution of the tissue-specific vs. tissue-shared m⁶A-QTLs in each tissue. **k.** Gene expression in tissues for tissue-specific gmGenes in each tissue. Statistical test was carried out by two-sided paired Wilcoxon test. **l.** Correlation (adjusted R²) of eQTLs between GTEx primary tissues and YRI LCL cells. Boxes=25%-75% percentile; line=median; whiskers=1.5 IQR. **m.** Comparison of m⁶A-QTL effects size between this study (eGTEx tissues, x-axis) and the other m⁶A-QTL study (YRI LCLs, y-axis). Green dots represent the m⁶A-QTLs shared by the two studies. Directionality consistency and corresponding p-value (vs. the 50% expected by chance) calculated using one-sided Fisher exact test (inset box). **n.** Correlation of m⁶A-QTLs between eGTEx primary tissues and YRI LCLs. **o.** Correlation between p-value and m⁶A-QTL effect in LCL cell lines for m⁶A-QTLs identified in eGTEx tissues, with positive- and negative-effect eGTEx m⁶A-QTLs separated. Directionality consistency and corresponding p-value (vs. the 50% expected by chance) calculated using one-sided Fisher exact test.



Extended Data Fig. 4 | Comparison between eQTL and m⁶A-QTL. **a-b.** m⁶A-QTLs show a modest but significant enrichment for eQTLs in the matching tissues, with m⁶A-QTLs separated into exonic (b) and intronic (c). NAnnot=21811, 17687, 40078, 24317, 39687 for Brain Cortex, Brain Frontal Cortex (BA9), Lung, Heart Left Ventricle, Skeletal Muscle (see Methods) for exonic m⁶A-QTLs. NAnnot=23091, 18724, 41807, 25927, 42425 for Brain Cortex, Brain Frontal Cortex (BA9), Lung, Heart Left Ventricle, Skeletal Muscle (see Methods) for intronic m⁶A-QTLs. Error bars denote the upper bound and the lower bound for the 95% CI of effect size. P-values are calculated by Garfield using a logistic regression model with ‘feature matching’. **c.** Number of eQTLs identified (y-axis) for increasing number of PEER factors removed (x-axis) shows inflection-point for each tissue (colors). **d.** Genomic region distribution for g-m⁶As mediating stabilization vs. degradation (p-values: Fisher exact test). **e.** Overlap between gmGenes and eGenes identified from the matching GTEx individuals. **f.** Effect size comparison between m⁶A-QTL and GTEx eQTL for the m⁶A-QTLs identified in this study. **g.** Effect size comparison between m⁶A-QTL and GTEx eQTL for the eQTLs identified by GTEx V8.

Extended Data Fig. 5 | Overlap between m⁶A-QTL and disease GWAS hits. **a-b.** Overlaid Manhattan plots showing genomic position (y-axis) and m⁶A-QTL P-value (x-axis) for lead SNPs (points) across traits (colors) that show colocalization between GWAS variants and m⁶A-QTLs in lung (a) and muscle/heart (b). **c.** Illustrative example showing a brain intronic m⁶A-QTL that is overlapped with a ClinVar-curated variant related to Congenital cataract. **d.** Illustrative example showing a muscle/heart 5'-UTR m⁶A-QTL that is overlapped with a ClinVar-curated variant related to Nemaline myopathy. Boxes = 25%-75% percentile; line=median; whiskers=1.5 IQR. **e.** Same plot as Fig. 5d, but shown using S-LDSC Baseline v1.2, which corrects for coding region, UTR, intron, promoter, enhancer, multiple histone marks, and eQTLs, shows robustness of results to this correction.



Extended Data Fig. 6 | Identification of m⁶A regulator candidates. **a.** Enrichment (y axis, log) and corresponding p-value (x-axis, Bonferroni-corrected) between m⁶A-QTL SNPs and RNA binding protein (RBP) binding sites, for each tissue (color), highlighting 10 most enriched RBPs in each tissue (labels). **b.** Quantile-Quantile plot showing the p-value distribution observed in the correlation test between RBP expression vs. m⁶A levels (y-axis), compared to the non-target (circle) or permutation controls (triangle) (x-axis). Significant RBPs (FDR < 0.1) are shown in red. P-values are calculated by two-sided Pearson correlation tests. **c,d.** Correlation of predicted m⁶A-regulator RBP mRNA expression level (x-axis) vs. methylation level of its m⁶A targets (y-axis) for RBM15 and FIP1L1 in brain, respectively. Grey shadow denotes the 95% confidence region for the regression fit.

Corresponding author(s): Manolis KellisLast updated by author(s): May 1, 2021

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection The m6A profiling sequencing data was generated at the Genomics Platform and the Broad Technology Labs (BTL) using the illumina Hiseq 4000 sequencer. Raw sequencing data was aligned and processed using bowtie2 (v2.3.4.3), hisat2 (v2.1.0) and samtools (v1.9).

Data analysis We used MACS2 (v2.1.1) for peak calling, bedtools (v2.28.0) for peak processing, deeptools (3.0.2) for coverage analyses, AME (v5.0.3) for motif enrichment analysis, R (v3.6.1) for all the statistic analysis, PEER (R package 1.0) for latent factor identification, FastQTL (v2.0) for QTL identification, GARFIELD (version 2) for localization enrichment analyses between m6A-QTL vs. RBP binding site and eQTL, LDSC (v1.0.1) for GWAS heritability analysis, STRINGdb (R package 1.24.0) for PPI analysis.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

All eGTEX protected data, including m6A sequencing reads and matched RNA-seq data are available on dbGaP with accession number phs000424.v8.p2. Additionally, the data can be accessed via AnVIL with authentication: https://anvil.terra.bio/#workspaces/anvil-datastorage/AnVIL_GTEX_V8_hg38. Since the raw sequencing data with human genetic information are protected, application and authentication are needed before accessing the data. All non-protected data of m6A can be visualized via the GTEX Portal (www.gtexportal.org) as part of eGTEX v8. The eQTL datasets are from GTEX v8, which can be accessed at <https://>

gtexportal.org/home/datasets. The LCL m6A-QTL datasets from Zhang et al. can be downloaded from <https://doi.org/10.5281/zenodo.3870952>. The previously curated m6A sites can be downloaded from RMBase (<http://rna.sysu.edu.cn/rmbase/>). The RNA binding sites can be downloaded from <http://lulab.life.tsinghua.edu.cn/postar/>.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	Due to the limited availability of human primary tissue samples, we initially performed 176 transcriptome-wide m6A profiling across 107 individuals in human brain, heart, muscle, and lung, and after stringent quality control, 129 m6A profiles were kept for analyses. Although the sample size may not be sufficient to identify a large amount of m6A-QTL, we believe that as a pioneer m6A genetics study in human tissues, these datasets already enable us to discover the m6A-QTL with strong effect for each of the tissues and to investigate many fundamental m6A genetic questions. No statistical methods were used to predetermine sample size.
Data exclusions	We excluded 47 m6A profiles prior to any m6A-QTL analysis, based on the m6A motif enrichment and the m6A peak distribution pattern. These are the two well-known metrics for m6A data quality evaluation, and such exclusion criteria were predefined based on the m6A profiling paper (Dan Dominissini et al., Nature 2012).
Replication	The experiments across 176 samples were independently carried out in three batches (see the Methods and Table S1 for details), among which 129 samples showed qualified m6A metrics. The rest of 47 samples failed possibly due to the RNA degradation of the biosamples.
Randomization	Randomization is not relevant to this m6A-QTL study, as the grouping of individuals was based on the genotype for the individuals in each locus, and can be varied across different locus.
Blinding	As described above, grouping of samples was naturally blinded as it is based on the genotypes of each individual in each locus.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

Methods

n/a	Involved in the study
<input type="checkbox"/>	<input checked="" type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Antibodies

Antibodies used	Anti-m6A polyclonal antibody (Synaptic Systems, Cat. no 202 003, Lot 202003/18), 1:200 diluted for IP usage.
Validation	Validated on human for western blotting, IP, ICC, IHC-P/FFPE and ELISA by SYSY. The antibody is against N6-methyladenosine modifications of RNA and DNA, reacting with: human, rat, mouse, eukaryotes, prokaryotes. Other species not tested yet. The detailed validation information of this antibody is available at: https://sysy.com/product/202003 .